

03/30/00
Jc644 U.S. PTO

Jc598 U.S. PTO
09/537710
03/30/00

Please type a plus sign (+) inside this box->/+ /

UTILITY	Atty Doc. No. 3377/99-Util	Total Page 59
PATENT APPLICATION	FIRST NAMED INVENTOR OR APPLICATION IDENTIFIER	
TRANSMITTAL	Anders DAHLQVIST	
	Express Mail Label No.	

Application Elements

Address To: Assistant Commissioner for Patents
Box Patent Application
Washington, D.C. 20231

1. / X / Fee transmittal Form
(Submit an original, and a duplicate for fee processing)
2. / X / Specification Total Pages /
(Preferred arrangement set for below)

6. / / Microfiche Computer Program (Appendix)
7. / / Nucleotide and/or Amino Acid Sequence Submission
(if applicable, all necessary)

Descriptive title of the Invention
Cross References to Related Application
Statement Regarding Fed. Sponsored R & D
Reference to Microfiche Appendix
Background of the Invention
Brief Summary of the Invention
Brief Description of the Drawings (if filed)
Detailed Description
Claim(s)
Abstract of the Disclosure

- a. / / Computer Readable Copy
b. / / Paper Copy (Identical to computer copy)
c. / / Statement verifying identity of above copies
ACCOMPANYING APPLICATIONS PARTS
8. / / Assignment Papers (cover sheet & document(s))
9. / 37 CFR 3.73(b) Statement / / Power of Attorney
10. / / English Translation Document (if applicable)
11. / / Information Disclosure / / Copies of IDS Citations
12. / / Preliminary Amendment
13. / x / Return Receipt Postcard (MPEP 503)

3. / X / Drawing(s) (35 USC 113) (Figs.) Total Sheets / 6 /
4. / / Oath or Declaration Total Pages / /

- Should be specifically itemized)
14. / / Small Entity / / Statement filed in prior application
Statements Status still proper and desired
15. / / Certified Copy of Priority Document(s)
(if foreign priority is claimed)
16. / / Other

- a. / / Newly executed (original or copy)
b. / / Copy from a prior application (37 CFR 1.63(d))
(For Continuation/Divisional with Box 17 completed)
Note Box 5 below
i. / / DELETION OF INVENTOR(S)
Signed statement attached deleting
inventor(s) named in the prior application
see 37 CFR 1.63(d)(2) and 1.33(b).

5. / / Incorporation by reference (useable if Box 4b is checked)
The entire disclosure of the prior application, from which a
copy of the oath or declaration is supplied under Box 4b
is considered as being part of the disclosure of the accompanying
application and is hereby incorporated by reference therein.

17. If a Continuing Application, check appropriate box and supply the requisite information:

/ X / Continuation / / Divisional / / Continuation-in part (CIP) of prior application No. 06/180,687

CORRESPONDENCE ADDRESS

/ / Customer Number or Bar code Label

or / / Correspondence address below

Insert Customer No. or Attach bar code label here

Name: Herbert B. Keil
KEIL & WEINKAUF

Address: 1101 Connecticut Ave., N.W.

City: Washington

State: D.C.

Zip Code 20036

Country: USA

Telephone: (202)659-0100

Fax: (202)659-0105

The filing fee has been calculated as shown below:

For:	Number Filed	Number Extra	SMALL/LARGE ENTITY	BASIC FEE \$345./\$690.
------	-----------------	-----------------	-----------------------	----------------------------

Basic Fee.....				\$ 690.00
Total Claims: 30	-20	=	x \$09./\$18.	= 180.00
Indep. Claims:	-3	=	x \$39./\$78.	=

[X] Multiple Dependent Claim(s) presented: \$130./260 = 260.00

[x] A check is enclosed for the filing fee. \$ 1,130.00

*If the difference is less than zero, enter "0".

[X] A check for \$1,130.00 for the filing fee and recordation fee.

[X] The Commissioner is hereby authorized to charge any other fee required, including the issue fee, in connection with the filing and prosecution of this application, and to the extent necessary, applicant(s) hereby petition for extension(s) of time under 37 CFR 1.136, to be charged to our Deposit Account 11-0345.

Respectfully submitted,
KEIL & WEINKAUF

H B Keil
Herbert B. Keil
Reg. No. 18,967

1101 Connecticut Ave., N.W
Washington, D.C. 20036
(202) 659-0100

000000-0744500

1	100%
2	98%
3	96%
4	94%
5	92%
6	90%
7	88%
8	86%
9	84%
10	82%
11	80%
12	78%
13	76%
14	74%
15	72%
16	70%
17	68%
18	66%
19	64%
20	62%
21	60%
22	58%
23	56%
24	54%
25	52%
26	50%
27	48%
28	46%
29	44%
30	42%
31	40%
32	38%
33	36%
34	34%
35	32%
36	30%
37	28%
38	26%
39	24%
40	22%
41	20%
42	18%
43	16%
44	14%
45	12%
46	10%
47	8%
48	6%
49	4%
50	2%
51	0%

)))))))

11

Group Art:

1

Examiner

1

For: A NEW CLASS OF ENZYMES IN THE BIOSYNTHETIC PATHWAY FOR THE PRODUCTION OF TRIACYLGLYCEROL AND RECOMBINANT DNA MOLECULES ENCODING THESE ENZYMES

;

1

7

E

○

7

1

1

A

•

de

Claim 12, lines 1 and 2, delete "claims 7 to 11" and insert --claim 7- -

Claim 13, lines 1 and 2, delete "claims 7 to 11 or a gene construct according to claim 12" and insert --claim 7--.

Claim 15, line 1, delete "claims 13 or 14" and insert --claim 13--.

Claim 16, line 2, delete "claims 7 to 11" and insert --claim 7- -;

line 3, delete "to 15" and insert --28--.

Claim 18, line 1, delete "claims 16 or 17" and insert --claim 16--.

Claim 19, line 1, delete "claims 16 to 18" and insert --claim 16--.

Claim 20, line 1, delete "claims 16 to 19" and insert --claim 19--.

Claim 21, line 1, delete "claims 16 to 20" and insert --claim 16--.

Claim 22, line 1, delete "claims 16 to 21" and insert --claim 16--.

Claim 23, line 1, delete "claims 16 to 22" and insert --claim 16--.

Claim 25, line 1, change "24" to ---30--.

Claim 15, line 1, delete "24" and insert --20--.

Cancel claims 24, 26 and 27.

Insert the following new claims.

--28. A vector comprising the gene construct of claim 12.

29. A vector according to claim 28, further comprising a selectable master gene and/or nucleotide sequences for the replication in a host cell or the integration into the genome of the host cell.

30. A process for the production of triacylglycerol, comprising growing a transgenic cell or organism according to claim 16 under conditions whereby a nucleotide sequence encoding an enzyme catalyzing in an acyl-CoA-independent reaction the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol

is expressed and whereby said transgenic cells comprising an enzyme catalyzing in a acyl-CoA-independent reaction the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol.

31. A method of producing triacylglycerol and/or triacylglycerol with uncommon fatty acids which comprises transforming an organism or host cell using the nucleotide sequence of claim 7, whereby the transformation results in an altered, preferably, increased oil content of the cell or organism.

32. A method of producing triacylglycerol and/or triacylglycerols with uncommon fatty acids using the nucleotide sequence of claim 7.

33. A method of producing triacylglycerol and/or triacylglycerols with uncommon fatty acids using the enzyme of claim 1.


REMARKS

The claims have been amended to put the application in better form for U.S. filing. No new matter has been added.

Entry of the above amendment is respectfully solicited.

Respectfully submitted,

KEIL & WEINKAUF


Herbert B. Keil
Reg. No. 18,967

1101 Connecticut Ave., N.W.
Washington, D.C. 20036
(202)659-0100

A NEW CLASS OF ENZYMES IN THE BIOSYNTHETIC PATHWAY FOR THE
PRODUCTION OF TRIACYLGLYCEROL AND RECOMBINANT DNA
MOLECULES ENCODING THESE ENZYMES

- 5 The present invention relates to the isolation, identification and characterization of recombinant DNA molecules encoding enzymes catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol.
- 10 Triacylglycerol (TAG) is the most common lipid-based energy reserve in nature. The main pathway for synthesis of TAG is believed to involve three sequential acyl-transfers from acyl-CoA to a glycerol backbone (1, 2). For many years, acyl-CoA : diacylglycerol acyltransferase (DAGAT), which catalyzes the third acyl transfer reaction, was thought to be the only unique enzyme involved in
- 15 TAG synthesis. It acts by diverting diacylglycerol (DAG) from membrane lipid synthesis into TAG (2). Genes encoding this enzyme were recently identified both in the mouse (3) and in plants (4, 5), and the encoded proteins were shown to be homologous to acyl-CoA : cholesterol acyltransferase (ACAT). It was also recently reported that another DAGAT exists in the oleaginous fungus
- 20 *Mortierella ramanniana*, which is unrelated to the mouse DAGAT, the ACAT gene family or to any other known gene (6).

The instant invention relates to novel type of enzymes and their encoding genes for transformation. More specifically, the invention relates to use of a

25 type of genes encoding a not previously described type of enzymes hereinafter designated phospholipid:diacylglycerol acyltransferases (PDAT), whereby this enzyme catalyses an acyl-CoA-independent reaction. The said type of genes expressed alone in transgenic organisms will enhance the total amount of oil (triacylglycerols) produced in the cells. The PDAT genes, in combination with a

30 gene for the synthesis of an uncommon fatty acid will, when expressed in transgenic organisms, enhance the levels of the uncommon fatty acids in the triacylglycerols.

There is considerable interest world-wide in producing chemical feedstock, such as fatty acids, for industrial use from renewable plant resources rather than non-renewable petrochemicals. This concept has broad appeal to
5 manufacturers and consumers on the basis of resource conservation and provides significant opportunity to develop new industrial crops for agriculture.

There is a diverse array of unusual fatty acids in oils from wild plant species and these have been well characterised. Many of these acids have industrial
10 potential and this has led to interest in domesticating relevant plant species to enable agricultural production of particular fatty acids.

Development in genetic engineering technologies combined with greater understanding of the biosynthesis of unusual fatty acids now makes it possible
15 to transfer genes coding for key enzymes involved in the synthesis of a particular fatty acid from a wild species into domesticated oilseed crops. In this way individual fatty acids can be produced in high purity and quantities at moderate costs.

20 In all crops like rape, sunflower, oilpalm etc., the oil (i.e. triacylglycerols) is the most valuable product of the seeds or fruits and other compounds like starch, protein, and fibre is regarded as by-products with less value. Enhancing the quantity of oil per weight basis at the expense of other compounds in oil crops would therefore increase the value of crop. If genes, regulating the allocation of
25 reduced carbon into the production of oil can be up-regulated, the cells will accumulate more oil on the expense of other products. Such genes might not only be used in already high oil producing cells such as oil crops but could also induce significant oil production in moderate or low oil containing crops such as e.g. soy, oat, maize, potato, sugarbeets, and turnips as well as in micro-
30 organisms.

Summary of the invention

Many of the unusual fatty acids of interest, e.g. medium chain fatty acids, hydroxy fatty acids, epoxy fatty acids and acetylenic fatty acids, have physical properties that are distinctly different from the common plant fatty acids. The present inventors have found that, in plant species naturally accumulating these uncommon fatty acids in their seed oil (i.e. triacylglycerol), these acids are absent, or present in very low amounts in the membrane (phospho)lipids of the seed. The low concentration of these acids in the membrane lipids is most likely a prerequisite for proper membrane function and thereby for proper cell functions. One aspect of the invention is that seeds of transgenic crops can be made to accumulate high amounts of uncommon fatty acids if these fatty acids are efficiently removed from the membrane lipids and channelled into seed triacylglycerols.

The inventors have identified a novel class of enzymes in plants catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the production of triacylglycerol through an acyl-CoA-independent reaction and that these enzymes (phospholipid:diacylglycerol acyltransferases abbreviated as PDAT) are involved in the removal of hydroxylated, epoxygenated fatty acids, and probably also other uncommon fatty acids such as medium chain fatty acids, from phospholipids in plants.

This enzyme reaction was shown to be present in microsomal preparations from baker's yeast (*Saccharomyces cerevisiae*). The instant invention further pertains to an enzyme comprising an amino acid sequence as set forth in SEQ ID No. 2 or a functional fragment, derivate, allele, homologue or isoenzyme thereof. A so called 'knock out' yeast mutant, disrupted in the respective gene was obtained and microsomal membranes from the mutant was shown to totally lack PDAT activity. Thus, it was proved that the disrupted gene encodes for a PDAT enzyme (SEQ ID NO. 1 and 2).

The instant invention pertains further to an enzyme comprising an amino acid sequence as set forth in SEQ ID NO. 1a, 2b or 5a or a functional fragment, derivate, allele, homologue or isoenzyme thereof.

Further genes and/or proteins of so far unknown function were identified and are contemplated within the scope of the instant invention. A gene from *Schizosaccharomyces pombe*, SPBC776.14 (SEQ ID NO. 3), a putative open reading frame CAA22887 of the SPBC776.14 (SEQ ID NO. 13) were identified. Further *Arabidopsis thaliana* genomic sequences (SEQ ID NO. 4, 10 and 11) coding for putative proteins were identified, as well as a putative open reading frame AAC80628 from the *A. thaliana* locus AC 004557 (SEQ ID NO. 14) and a putative open reading frame AAD10668 from the *A. thaliana* locus AC 003027 (SEQ ID NO. 15) were identified.

Also, a partially sequenced cDNA clone from *Neurospora crassa* (SEQ ID NO. 9) and a *Zea mays* EST (Extended Sequence Tag) clone (SEQ ID NO. 7) and corresponding putative amino acid sequence (SEQ ID NO. 8) were identified. Finally, two cDNA clones were identified, one *Arabidopsis thaliana* EST (SEQ ID NO. 5 and corresponding predicted amino acid sequence SEQ ID NO. 6) and a *Lycopersicon esculentum* EST clone (SEQ ID NO. 12) were identified. Further, enzymes designated as PDAT comprising an amino acid sequence selected from the group consisting of sequences as set forth in SEQ ID NO 2a, 3a, 5b, 6 or 7b are contemplated within the scope of the invention. Moreover, an enzyme comprising an amino acid sequence encoded through a nucleotide sequence, a portion, derivate, allele or homologue thereof selected from the group consisting of sequences as set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 5, 5b, 6b, 7, 8b, 9, 9b, 10, 10b, 11, 11b or 12 or a functional fragment, derivate, allele, homologue or isoenzyme of the enzyme encoding amino acid sequence are included within the scope of the invention.

A functional fragment of the instant enzyme is understood to be any polypeptide sequence which shows specific enzyme activity of a phospholipid:diacylglycerol acyltransferase (PDAT). The length of the functional fragment can for example

vary in a range from about 660 ± 10 amino acids to 660 ± 250 amino acids, preferably from about 660 ± 50 to 660 ± 100 amino acids, whereby the „basic number“ of 660 amino acids corresponds in this case to the polypeptide chain of the PDAT enzyme of SEQ ID NO. 2 encoded by a nucleotide sequence according to SEQ ID NO. 1. Consequently, the „basic number“ of functional fulllength enzyme can vary in correspondance to the encoding nucleotide sequence.

A portion of the instant nucleotide sequence is meant to be any nucleotide sequence encoding a polypeptid which shows specific activity of a phospholipid:diacylglycerol acyltransferase (PDAT). The length of the nucleotide portion can vary in a wide range of about several hundereds of nucleotides based upon the coding region of the gene or a highly conserved sequence. For example the length varies in a range form about 1900 ± 10 to 1900 ± 1000 nucleotides, preferably form about 1900 ± 50 to 1900 ± 700 and more preferably form about 1900 ± 100 to 1900 ± 500 nucleotides. whereby the „basic number“ of 1900 nucleotides corresponds in this case to the encoding nucleotide sequence of the PDAT enzyme of SEQ ID NO. 1. Consequently, the „basic number“ of functional fulllength gene can vary.

An allelic variant of the instant nucleotide sequence is understood to be any different nucleotide sequence which encodes a polypeptide with a functionally equivalent function. The alleles pertain naturally occuring variantes of the instant nucleotide sequences as well as synthetic nucleotide sequences produces by methods known in the art. Contemplated are even altered nucleotide sequences which result in an enzyme with altered activity and/or regulation or which is resistant against specific inhibitors. The instant invention further includes natural or synthetic mutations of the originally isolated nucleotide sequences. These mutations can be substitution, addition, deletion, inversion or insertion of one or more nucleotides.

A homologues nucleotide sequence is understood to be a complementary sequence and/or a sequence which specifically hybridizes with the instant nucleotide sequence. Hybridizing sequences include similar sequences selected from the group of DNA or RNA which specifically interact to the instant nucleotide sequences under at least moderate stringency conditions which are known in the art. A preferred, non-limiting example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. This further includes short nucleotide sequences of e.g. 10 to 30 nucleotides, preferably 12 to 15 nucleotides. Included are also primer or hybridization probes.

A homologue nucleotide sequence included within the scope of the instant invention is a sequence which is at least about 40%, preferably at least about 50 % or 60%, and more preferably at least about 70%, 80% or 90% and most preferably at least about 95%, 96%, 97%, 98% or 99% or more homologous to a nucleotide sequence of SEQ ID NO. 1.

All of the aforementioned definitions are true for amino acid sequences and functional enzymes and can easily transferred by a person skilled in the art.

Isoenzymes are understood to be enzymes which have the same or a similar substrate specificity and/or catalytic activity but a different primary structure.

In a first embodiment, this invention is directed to nucleic acid sequences that encode a PDAT. This includes sequences that encode biologically active PDATs as well as sequences that are to be used as probes, vectors for transformation or cloning intermediates. The PDAT encoding sequence may encode a complete or partial sequence depending upon the intended use. All or a portion of the genomic sequence, cDNA sequence, precursor PDAT or mature PDAT is intended.

Further included is a nucleotide sequence selected from the group consisting of sequences set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 9b, 10, 10b or 11 or a portion, derivate, allele or homologue thereof. The invention pertains a partial
5 nucleotide sequence corresponding to a fulllength nucleotide sequence selected from the group consisting of sequences set forth in SEQ ID No. 5, 5b, 6b, 7, 8b, 9, 11b or 12 or a portion, derivate, allele or homologue thereof. Moreover, a nucleotide sequence comprising a nucleotide sequence which is at least 40% homologous to a nucleotide sequence selected form the group consisting of
10 those sequences set forth in SEQ ID No. 1 1b, 3, 3b, 4, 4a, 4b, 5, 5b, 6b, 7, 8b, 9, 9b, 10, 10b, 11, 11b or 12 is contemplated within the scope of the invention.

The instant invention pertains to a gene construct comprising a said nucleotide sequences of the instant invention which is operably linked to a heterologous
15 nucleic acid.

The term operably linked means a serial organisation e.g. of a promotor, coding sequence, terminator and/or further regulatory elements whereby each element can fulfill its original function during expression of the nucleotide sequence.

20 Further, a vector comprising of a said nucleotide sequence of the instant invention is contemplated in the instant invention. This includes also an expression vector as well as a vector further comprising a selectable marker gene and/or nucleotide sequences for the replication in a host cell and/or the integration into the genome of the host cell.

25

In a different aspect, this invention relates to a method for producing a PDAT in a host cell or progeny thereof, including genetically engineered oil seeds, yeast and moulds or any other oil accumulating organism, via the expression of a construct in the cell. Cells containing a PDAT as a result of the production of
30 the PDAT encoding sequence are also contemplated within the scope of the invention.

Further, the invention pertains a transgenic cell or organism containing a said nucleotide sequence and/or a said gene construct and/or a said vector. The object of the instant invention is further a transgenic cell or organism which is an eucaryotic cell or organism. Preferably, the transgenic cell or organism is a yeast cell or a plant cell or a plant. The instant invention further pertains said transgenic cell or organism having an altered biosynthetic pathway for the production of triacylglycerol. A transgenic cell or organism having an altered oil content is also contemplated within the scope of this invention.

10

Further, the invention pertains a transgenic cell or organism wherein the activity of PDAT is altered in said cell or organism. This altered activity of PDAT is characterized by an alteration in gene expression, catalytic activity and/or regulation of activity of the enzyme. Moreover, a transgenic cell or organism is included in the instant invention, wherein the altered biosynthetic pathway for the production of triacylglycerol is characterized by the prevention of accumulation of undesirable fatty acids in the membrane lipids.

15

In a different embodiment, this invention also relates to methods of using a DNA sequence encoding a PDAT for increasing the oil-content within a cell.

20

Another aspect of the invention relates to the accommodation of high amounts of uncommon fatty acids in the triacylglycerol produced within a cell, by introducing a DNA sequence producing a PDAT that specifically removes these fatty acids from the membrane lipids of the cell and channel them into triacylglycerol. Plant cells having such a modification are also contemplated herein.

25

Further, the invention pertains a process for the production of triacylglycerol, comprising growing a said transgenic cell or organism under conditions whereby the said nucleotide sequence is expressed and whereby the said

30

transgenic cells comprising an said enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol forming triacylglycerol.

Moreover, triacylglycerols produced by the aforementioned process are
5 included in scope of the instant invention.

Object of the instant invention is further the use of an instant nucleotide sequence and/or a said enzyme for the production of triacylglycerol and/or triacylglycerols with uncommon fatty acids. The use of a said instant nucleotide
10 sequence and/or a said enzyme of the instant invention for the transformation of any cell or organism in order to be expressed in this cell or organism and result in an altered, preferably increased oil content of this cell or organism is also contemplated within the scope of the instant invention.

15 A PDAT of this invention includes any sequence of amino acids, such as a protein, polypeptide or peptide fragment obtainable from a microorganism, animal or plant source that demonstrates the ability to catalyse the production of triacylglycerol from a phospholipid and diacylglycerol under enzyme reactive conditions. By „enzyme reactive conditions,, is meant that any necessary
20 conditions are available in an environment (e.g., such factors as temperature, pH, lack of inhibiting substances) which will permit the enzyme to function.

Other PDATs are obtainable from the specific sequences provided herein. Furthermore, it will be apparent that one can obtain natural and synthetic
25 PDATs, including modified amino acid sequences and starting materials for synthetic-protein modelling from the exemplified PDATs and from PDATs which are obtained through the use of such exemplified sequences. Modified amino acid sequences include sequences that have been mutated, truncated, increased and the like, whether such sequences were partially or wholly
30 synthesised. Sequences that are actually purified from plant preparations or are identical or encode identical proteins thereto, regardless of the method

used to obtain the protein or sequence, are equally considered naturally derived.

Further, the nucleic acid probes (DNA and RNA) of the present invention can
5 be used to screen and recover „homologous,, or „related,, PDATs from a variety of plant and microbial sources.

Further, it is also apparent that a person skilled in the art can, with the information provided in this application, in any organism identify a PDAT
10 activity, purify an enzyme with this activity and thereby identify a „non-homologues“ nucleic acid sequence encoding such an enzyme.

The present invention can be essentially characterized by the following aspects:

15

1. Use of a PDAT gene (genomic clone or cDNA) for transformation.
2. Use of a DNA molecule according to item 1 wherein said DNA is used for transformation of any organism in order to be expressed in this organism and result in an active recombinant PDAT enzyme in order to increase oil
20 content of the organism.
3. Use of a DNA molecule of item 1 wherein said DNA is used for transformation of any organism in order to prevent the accumulation of undesirable fatty acids in the membrane lipids.
4. Use according to item 1, wherein said PDAT gene is used for transforming
25 transgenic oil accumulating organisms engineered to produce any uncommon fatty acid which is harmful if present in high amounts in membrane lipids, such as medium chain fatty acids, hydroxylated fatty acids, epoxygenated fatty acids and acetylenic fatty acids.
5. Use according to item 1, wherein said PDAT gene is used for transforming
30 organisms, and wherein said organisms are crossed with other oil accumulating organisms engineered to produce any uncommon fatty acid

which is harmful if present in high amounts in membrane lipids, comprising medium chain fatty acids, hydroxylated fatty acids, epoxygenated fatty acids and acetylenic fatty acids.

6. Use according to item 1, wherein the enzyme encoded by said PDAT gene or cDNA is coding for a PDAT with distinct acyl specificity.
7. Use according to item 1 wherein said PDAT encoding gene or cDNA, is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 30% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
8. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 40% or more *identical* to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
9. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 60% or more *identical* to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
10. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 80% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
11. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from plants or contain nucleotide sequences coding for an amino acid sequence 40% or more identical to the amino acid sequence of PDAT from *Arabidopsis thaliana* or to the protein encoded by the fulllength counterpart of the partial *Zea mays*, *Lycopersicon esculentum*, or *Neurospora crassa* cDNA clones.
12. Transgenic oil accumulating organisms comprising, in their genome, a PDAT gene transferred by recombinant DNA technology or somatic hybridization.

13. Transgenic oil accumulating organisms according to item 12 comprising, in their genome, a PDAT gene having specificity for substrates with a particular uncommon fatty acid and the gene for said uncommon fatty acid.
14. Transgenic organisms according to item 12 or 13 which are selected from the group consisting of fungi, plants and animals.
15. Transgenic organisms according to item 12 or 13 which are selected from the group of agricultural plants.
16. Transgenic organisms according to item 12 or 13 which are selected from the group of agricultural plants and where said PDAT gene is expressed under the control of a storage organ specific promotor.
17. Transgenic organisms according to item 12 or 13 which are selected from the group of agricultural plants and where said PDAT gene is expressed under the control of a seed promotor.
18. Oils from organisms according to item 12 – 17.
19. A method for altering acyl specificity of a PDAT by alteration of the nucleotide sequence of a naturally occurring encoding gene and as a consequence of this alternation creating a gene encoding for an enzyme with novel acyl specificity.
20. A protein encoded by a DNA molecule according to item 1 or a functional fragment thereof.
21. A protein of item 20 designated phospholipid:diacylglycerol acyltransferase.
22. A protein of item 21 which has a distinct acyl specificity.
23. A protein of item 13 having the amino acid sequence as set forth in SEQ, ID NO. 2, 13, 14 or 15 (and the proteins encoded by the fulllength or partial genes set forth in SEQ. ID. NO. 1, 3, 4, 5, 7, 9, 10, 11 or 12) or an amino acid sequence with at least 30 % homology to said amino acid sequence.
24. A protein of item 23 isolated from *Saccharomyces cerevisiae*.

General methods:

Yeast strains and plasmids. The wild type yeast strains used were either FY1679 (*MAT α his3- Δ 200 leu2- Δ 1 trp1- Δ 6 ura3-52*) or W303-1A (*MAT α ADE2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1*) (7). The YNR008w::KanMX2 disruption strain FVKT004-04C(AL), which is congenic to FY1679, was
 5 obtained from the Euroscarf collection (8). A 2751 bp fragment containing the YNR008w gene with 583 bp of 5' and 183 bp of 3' flanking DNA was amplified from W303-1A genomic DNA using *Taq* polymerase with 5'-TCTCCATCTTCTGCAAAACCT-3' and 5'-CCTGTCAAAAACCTTCTCCTC-3' as
 10 primers. The resulting PCR product was purified by agarose gel electrophoresis and cloned into the *EcoRV* site of pBluescript (pbluescript-pdat). For complementation experiments, the cloned fragment was released from pBluescript by *HindIII*-*SacI* digestion and then cloned between the *HindIII* and *SacI* sites of pFL39 (9), thus generating pUS1. For overexpression of the PDAT gene, a 2202 bp *EcoRI* fragment from the pBluscript plasmid which contains
 15 only 24 bp of 5' flanking DNA was cloned into the *BamHI* site of the *GAL1*-*TPK2* expression vector pJN92 (12), thus generating pUS4.

Microsomal preparations. Microsomes from developing seeds of sunflower (*Helianthus annuus*), *Ricinus communis* and *Crepis palaestina* were prepared
 20 using the procedure of Stobart and Stymne (11). To obtain yeast microsomes, 1g of yeast cells (fresh weight) was re-suspended in 8 ml of ice-cold buffer (20 mM Tris-Cl, pH 7.9, 10 mM MgCl₂, 1 mM EDTA, 5 % (v/v) glycerol, 1 mM DTT, 0.3 M ammonium sulfate) in a 12 ml glass tube. To this tube, 4 ml of glass beads (diameter 0.45-0.5 mm) were added, and the tube was then heavily
 25 shaken (3 x 60 s) in an MSK cell homogenizer (B. Braun Melsungen AG, Germany). The homogenized suspension was centrifuged at 20,000 x g for 15 min at 6°C and the resulting supernatant was again centrifuged at 100,000 x g for 2 h at 6°C. The 100,000 x g pellet was resuspended in 0.1 M potassium phosphate (pH 7.2), and stored at -80°C. It is subsequently referred to as the
 30 crude yeast microsomal fraction.

Lipid substrates. Radio-labeled ricinoleic (12-hydroxy-octadecenoic) and vernolic (12,13-epoxy-octadecenoic) acids were synthesized enzymatically from [1-¹⁴C]oleic acid and [1-¹⁴C]linoleic acid, respectively, by incubation with
 5 microsomal preparations from seeds of *Ricinus communis* and *Crepis palaestina*, respectively (12). The synthesis of phosphatidylcholines (PC) or phosphatidylethanolamines (PE) with ¹⁴C-labeled acyl groups in the *sn*-2 position was performed using either enzymatic (13), or synthetic (14) acylation of [¹⁴C]oleic, [¹⁴C]ricinoleic, or [¹⁴C]vernolic acid. Dioleoyl-PC that was labeled
 10 in the *sn*-1 position was synthesized from *sn*-1-[¹⁴C]oleoyl-lyso-PC and unlabeled oleic acid as described in (14). *Sn*-1-oleoyl-*sn*-2-[¹⁴C]ricinoleoyl-DAG was synthesized from PC by the action of phospholipase C type XI from *B. Cereus* (Sigma Chemical Co.) as described in (15). Monovernoloyl- and divernoloyl-DAG were synthesized from TAG extracted from seeds of
 15 *Euphorbia lagascae*, using the TAG-lipase (*Rizhopus arrizus*, Sigma Chemical Co.) as previously described (16). Monoricinoleoyl-TAG was synthesized according to the same method using TAG extracted from Castor bean.

Lipid analysis. Total lipid composition of yeast were determined from cells
 20 harvested from a 40 ml liquid culture, broken in a glass-bead shaker and extracted into chloroform as described by Bligh and Dyer (17), and then separated by thin layer chromatography in hexane/diethylether/acetic acid (80:20:1) using pre-coated silica gel 60 plates (Merck). The lipid areas were located by brief exposure to I₂ vapors and identified by means of appropriate
 25 standards. Polar lipids, sterol-esters and triacylglycerols, as well as the remaining minor lipid classes, referred to as other lipids, were excised from the plates. Fatty acid methylesters were prepared by heating the dry excised material at 85 °C for 60 min in 2% (v/v) sulfuric acid in dry methanol. The methyl esters were extracted with hexane and analyzed by GLC through a 50 m
 30 x 0.32 mm CP-Wax58-CB fused-silica column (Chrompack), with methylheptadecanoic acid as an internal standard. The fatty acid content of

each fraction was quantified and used to calculate the relative amount of each lipid class. In order to determine the total lipid content, 3 ml aliquots from yeast cultures were harvested by centrifugation and the resulting pellets were washed with distilled water and lyophilized. The weight of the dried cells was determined and the fatty acid content was quantified by GLC-analyses after conversion to methylesters as described above. The lipid content was then calculated as nmol fatty acid (FA) per mg dry weight yeast.

Enzyme assays. Aliquots of crude microsomal fractions (corresponding to 10 nmol of microsomal PC) from developing plant seeds or yeast cells were lyophilized over night. ^{14}C -Labeled substrate lipids dissolved in benzene were then added to the dried microsomes. The benzene was evaporated under a stream of N_2 , leaving the lipids in direct contact with the membranes, and 0.1 ml of 50 mM potassium phosphate (pH 7.2) was added. The suspension was thoroughly mixed and incubated at 30°C for the time period indicated, up to 90 min. Lipids were extracted from the reaction mixture using chloroform and separated by thin layer chromatography in hexane/diethylether/acetic acid (35:70:1.5) using silica gel 60 plates (Merck). The radioactive lipids were visualized and quantified on the plates by electronic autoradiography (Instant Imager, Packard, US).

Yeast cultivation. Yeast cells were grown at 28°C on a rotatory shaker in liquid YPD medium (1% yeast extract, 2% peptone, 2% glucose), synthetic medium (18) containing 2% (v/v) glycerol and 2% (v/v) ethanol, or minimal medium (19) containing 16 g/l of glycerol.

The instant invention is further characterized by the following examples which are not limiting:

Acyl-CoA-independent synthesis of TAG by oil seed microsomes. A large number of unusual fatty acids can be found in oil seeds (20). Many of these

fatty acids, such as ricinoleic (21) and vernolic acids (22), are synthesized using phosphatidylcholin (PC) with oleoyl or linoleoyl groups esterified to the *sn*-2 position, respectively, as the immediate precursor. However, even though PC can be a substrate for unusual fatty acid synthesis and is the major membrane lipids in seeds, unusual fatty acids are rarely found in the membranes. Instead, they are mainly incorporated into the TAG. A mechanism for efficient and selective transfer of these unusual acyl groups from PC into TAG must therefore exist in oil seeds that accumulate such unusual fatty acids. This transfer reaction was biochemically characterized in seeds from castor bean (*Ricinus communis*) and *Crepis palaestina*, plants which accumulate high levels of ricinoleic and vernolic acid, respectively, and sunflower (*Helianthus annuus*), a plant which has only common fatty acids in its seed oil. Crude microsomal fractions from developing seeds were incubated with PC having ¹⁴C-labeled oleoyl, ricinoleoyl or vernoloyl groups at the *sn*-2 position. After the incubation, lipids were extracted and analyzed by thin layer chromatography. We found that the amount of radioactivity that was incorporated into the neutral lipid fraction increased linearly over a period of 4 hours (data not shown). The distribution of [¹⁴C]acyl groups within the neutral lipid fraction was analyzed after 80 min (Fig. 1). Interestingly the amount and distribution of radioactivity between different neutral lipids were strongly dependent both on the plant species and on the type of [¹⁴C]acyl chain. Thus, sunflower microsomes incorporated most of the label into DAG, regardless of the type of [¹⁴C]acyl group. In contrast, *R. communis* microsomes preferentially incorporated [¹⁴C]ricinoleoyl and [¹⁴C]vernoloyl groups into TAG, while [¹⁴C]oleyl groups mostly were found in DAG. *C. palaestina* microsomes, finally, incorporated only [¹⁴C]vernoloyl groups into TAG, with [¹⁴C]ricinoleyl groups being found mostly as free fatty acids, and [¹⁴C]oleyl groups in DAG. This shows that the high *in vivo* levels of ricinoleic acid and vernolic acid in the TAG pool of *R. communis* and *C. palaestina*, respectively, can be explained by an efficient and selective transfer of the corresponding acyl groups from PC to TAG in these organisms.

The in-vitro synthesis of triacylglycerols in microsomal preparations of developing castor bean is summarized in table 1.

PDAT: a novel enzyme that catalyzes acyl-CoA independent synthesis of

5 TAG. It was investigated if DAG could serve both as an acyl donor as well as an acyl acceptor in the reactions catalyzed by the oil seed microsomes. Therefore, unlabeled divernoloyl-DAG was incubated with either *sn*-1-oleoyl-*sn*-2-[¹⁴C]ricinoleoyl-DAG or *sn*-1-oleoyl-*sn*-2-[¹⁴C]ricinoleoyl-PC in the presence of *R. communis* microsomes. The synthesis of TAG molecules
10 containing both [¹⁴C]ricinoleoyl and vernoloyl groups was 5 fold higher when [¹⁴C]ricinoleoyl-PC served as acyl donor as compared to [¹⁴C]ricinoleoyl-DAG (fig.1B). These data strongly suggests that PC is the immediate acyl donor and DAG the acyl acceptor in the acyl-CoA-independent formation of TAG by oil seed microsomes. Therefore, this reaction is catalyzed by a new enzyme which
15 we call phospholipid : diacylglycerol acyltransferase (PDAT).

PDAT activity in yeast microsomes. Wild type yeast cells were cultivated under conditions where TAG synthesis is induced. Microsomal membranes were prepared from these cells and incubated with *sn*-2-[¹⁴C]-ricinoleoyl-PC and DAG and the ¹⁴C-labeled products formed were analyzed. The PC-derived
20 [¹⁴C]ricinoleoyl groups within the neutral lipid fraction mainly were found in free fatty acids or TAG, and also that the amount of TAG synthesized was dependent on the amount of DAG that was added to the reaction (Fig.2). The *in vitro* synthesis of TAG containing both ricinoleoyl and vernoloyl groups, a TAG
25 species not present *in vivo*, from exogenous added *sn*-2-[¹⁴C]ricinoleoyl-PC and unlabelled vernoloyl-DAG (Fig. 2, lane 3) clearly demonstrates the existence of an acyl-CoA-independent synthesis of TAG involving PC and DAG as substrates in yeast microsomal membranes. Consequently, TAG synthesis in yeast can be catalyzed by an enzyme similar to the PDAT found in plants.

The PDAT encoding gene in yeast.

A gene in the yeast genome (YNR008w) is known, but nothing is known about the function of YNR008w, except that the gene is not essential for growth under normal circumstances. Microsomal membranes were prepared from the yeast strain FVKT004-04C(AL) (8) in which this gene with unknown function had been disrupted. PDAT activity in the microsomes were assayed using PC with radiolabelled fatty acids at the sn-2 position. The activity was found to be completely absent in the disruption strain (Fig. 2 lane 4). Significantly, the activity could be partially restored by the presence of YNR008w on the single copy plasmid pUS1 (Fig. 2 lane 5). Moreover, acyl groups of phosphatidylethanolamine (PE) were efficiently incorporated into TAG by microsomes from the wild type strain whereas no incorporation occurred from this substrate in the mutant strain. This shows that YNR008w encodes a yeast PDAT which catalyzes the transfer of an acyl group from the sn-2 position of phospholipids to DAG, thus forming TAG. It should be noted that no cholesterol esters were formed from radioactive PC even in incubations with added ergosterols, nor were the amount of radioactive free fatty acids formed from PC affected by disruption of the YNR008w gene. This demonstrates that yeast PDAT do not have cholesterol ester synthesising or phospholipase activities.

Increased TAG content in yeast cells that overexpress PDAT. The effect of overexpressing the PDAT-encoding gene was studied by transforming a wild type yeast strain with the pUS4 plasmid in which the gene is expressed from the galactose-induced *GAL1:TPK2* promoter. Cells containing the empty expression vector were used as a control. The cells were grown in synthetic glycerol-ethanol medium, and expression of the gene was induced after either 2 hours (early log phase) or 25 hours (stationary phase) by the addition of galactose. The cells were then incubated for another 21 hours, after which they were harvested and assays were performed. We found that overexpression of PDAT had no significant effect on the growth rate as determined by the optical density. However, the total lipid content, measured as total μmol fatty acids per mg yeast dry weight, was 47% (log phase) or 29% (stationary phase) higher in the PDAT overexpressing strain than in the control. Furthermore, the polar lipid and sterolester content was unaffected by overexpression of PDAT. Instead, the elevated lipid content in these cells is entirely due to an increased TAG content (Fig. 3A,B). Thus, the amount of TAG was increased by 2-fold in PDAT overexpressing early log phase cells and by 40% in stationary phase cells. It is interesting to note that a significant increase in the TAG content was achieved by overexpressing PDAT even under conditions (*i.e.* in stationary phase) where DAGAT is induced and thus contributes significantly to TAG synthesis. *In vitro* PDAT activity assayed in microsomes from the PDAT overexpressing strain was 7-fold higher than in the control strain, a finding which is consistent with the increased levels of TAG that we observed *in vivo* (Fig. 3C). These results clearly demonstrate the potential use of the PDAT gene in increasing the oil content in transgenic organisms.

Substrate specificity of yeast PDAT. The substrate specificity of yeast PDAT was analyzed using microsomes prepared from the PDAT overexpressing strain (see Fig. 4). The rate of TAG synthesis, under conditions given in figure 4 with di-oleoyl-PC as the acyl-donor, was 0.15 nmol per min

and mg protein. With both oleoyl groups of PC labeled it was possible, under the given assay conditions, to detect the transfer of 11 pmol/min of [^{14}C]oleoyl chain into TAG and the formation of 15 pmol/min of lyso-PC. In microsomes from the PDAT-deficient strain, no TAG at all and only trace amounts of lyso-PC was detected, strongly suggesting that yeast PDAT catalyses the formation of equimolar amounts of TAG and lyso-PC when supplied with PC and DAG as substrates. The fact that somewhat more lyso-PC than TAG is formed can be explained by the presence of a phospholipase in yeast microsomes, which produces lyso-PC and unesterified fatty acids from PC (data not shown).

The specificity of yeast PDAT for different acyl group positions was investigated by incubating the microsomes with di-oleoyl-PC carrying a [^{14}C]acyl group either at the *sn*-1 position (Fig. 4A bar 2) or the *sn*-2 position (Fig. 4A bar 3). We found that the major ^{14}C -labeled product formed in the former case was lyso-PC, and in the latter case TAG. We conclude that yeast PDAT has a specificity for the transfer of acyl groups from the *sn*-2 position of the phospholipid to DAG, thus forming *sn*-1-lyso-PC and TAG. Under the given assay conditions, trace amounts of ^{14}C -labelled DAG is formed from the *sn*-1 labeled PC by the reversible action of a CDP-choline : choline phosphotransferase (data not shown). This labeled DAG can then be further converted into TAG by the PDAT activity. It is therefore not possible to distinguish whether the minor amounts of labeled TAG that is formed in the presence of di-oleoyl-PC carrying a [^{14}C]acyl group in the *sn*-1 position, is synthesized directly from the *sn*-1-labeled PC by a PDAT that also can act on the *sn*-1 position, or if it is first converted to *sn*-1-labeled DAG and then acylated by a PDAT with strict selectivity for the transfer of acyl groups at the *sn*-2 position of PC. Taken together, this shows that the PDAT encoded by YNR008w catalyses an acyl transfer from the *sn*-2 position of PC to DAG, thus causing the formation of TAG and lyso-PC.

The substrate specificity of yeast PDAT was further analyzed with respect to the headgroup of the acyl donor, the acyl group transferred and the acyl chains of the acceptor DAG molecule. The two major membrane lipids of *S. cerevisiae* are PC and PE, and as shown in Fig. 4B (bars 1 and 2), dioleoyl-PE is nearly 4-fold more efficient than dioleoyl-PC as acyl donor in the PDAT-catalyzed reaction. Moreover, the rate of acyl transfer is strongly dependent on the type of acyl group that is transferred. Thus, a ricinoleoyl group at the *sn*-2 position of PC is 2.5 times more efficiently transferred into TAG than an oleoyl group in the same position (Fig. 4B bars 1 and 3). In contrast, yeast PDAT has no preference for the transfer of vernoloyl groups over oleoyl groups (Fig. 4B bars 1 and 4). The acyl chain of the acceptor DAG molecule also affects the efficiency of the reaction. Thus, DAG with a ricinoleoyl or a vernoloyl group is a more efficient acyl acceptor than dioleoyl-DAG (Fig. 4B bars 1, 5 and 6). Taken together, these results clearly show that the efficiency of the PDAT-catalyzed acyl transfer is strongly dependent on the properties of the substrate lipids.

PDAT genes. Nucleotide and amino acid sequences of several PDAT genes are given as SEQ ID No. 1 through 15. Further provisional and/or partial sequences are given as SEQ ID NO 1a through 5a and 1b through 11b, respectively. One of the Arabidopsis genomic sequences (SEQ ID NO. 4) identified an Arabidopsis EST cDNA clone; T04806. This cDNA clone was fully characterised and the nucleotide sequence is given as SEQ ID NO. 5. Based on the sequence homology of the T04806 cDNA and the *Arabidopsis thaliana* genomic DNA sequence (SEQ ID NO 4) it is apparent that an additional A is present at position 417 in the cDNA clone (data not shown). Excluding this nucleotide would give the amino acid sequence depicted in SEQ ID NO. 12.

Increased TAG content in seeds of Arabidopsis thaliana that express the yeast PDAT. For the expression of the yeast pdat gene in *Arabidopsis thaliana* an EcoRI fragment from the pBluescript-pdat was cloned together with napin promoter (26) into the vector pGPTV-KAN (27). A plasmid (pGNapPDAT)

having the yeast PDAT gene in the correct orientation was identified and transformed into *Agrobacterium tumefaciens*. These bacteria were used to transform *Arabidopsis thaliana* columbia (C-24) plants using the root transformation method (28). Plants transformed with an empty vector were
5 used as controls.

First generation seeds (T1) were harvested and germinated on kanamycin containing medium. Second generation seeds (T2) were pooled from individual plants and their fatty acid contents analysed by quantification of their methyl esters by gas liquid chromatography after methylation of the seeds with 2%
10 sulphuric acid in methanol at 85 °C for 1,5 hours. Quantification was done with heptadecanoic acid methyl esters as internal standard.

From the transformation with pGNapPDAT one T1 plant (26-14) gave raise to seven T2 plants of which 3 plants yielded seeds with statistically (in a mean difference two-sided test) higher oil content than seeds from T2 plants
15 generated from T1 plant 32-4 transformed with an empty vector (table 2).

Claims

1. An enzyme catalysing in an acyl-CoA-independent reaction the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol.
2. An enzyme according to claim 1, comprising an amino acid sequence as set forth in SEQ ID No. 2 or a functional fragment, derivate, allele, homologue or isoenzyme thereof.
3. An enzyme according to claims 1 or 2 designated as phospholipid:diacylglycerol acyltransferase (PDAT).
4. An enzyme according to claims 1 to 3, comprising an amino acid sequence as set forth in SEQ ID No. 1a, 2b or 5a or a functional fragment, derivate, allele, homologue or isoenzyme thereof.
5. An enzyme according to claims 1 to 4, comprising an amino acid sequence selected from the group consisting of sequences as set forth in SEQ ID No. 2a, 3a, 5b, 6, 7b, 8, 13, 14, 15 or a functional fragment, derivate, allele, homologue or isoenzyme thereof.
6. An enzyme according to claims 1 to 5, comprising an amino acid sequence encoded through a nucleotide sequence, a portion, derivate, allele or homologue thereof selected from the group consisting of sequences as set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 5, 5b, 6b, 7, 8b, 9, 9b, 10, 10b, 11, 11b, 12 or a functional fragment, derivate, allele, homologue or isoenzyme of the enzyme encoding amino acid sequence.
7. A nucleotide sequence encoding an enzyme catalysing in an acyl-CoA-independent reaction the transfer of fatty acids from phospholipids to

diacylglycerol in the biosynthetic pathway for the production of triacylglycerol.

- 5
8. A nucleotide sequence according to claim 7 encoding an enzyme designated as phospholipid:diacylglycerol acyltransferase (PDAT).
9. A nucleotide sequence according to claims 7 or 8, selected from the group consisting of sequences as set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 9b, 10, 10b or 11 or a portion, derivate, allele or homologue thereof.
- 10
10. A partial nucleotide sequence corresponding to a fulllength nucleotide sequence according to claims 7 to 9, selected from the group consisting of sequences as set forth in SEQ ID No. 5, 5b, 6b, 7, 8b, 9, 11b or 12 or a portion, derivate, allele or homologue thereof.
- 15
11. A nucleotide sequence according to claims 7 to 10, comprising a nucleotide sequence which is at least 40% homologous to a nucleotide sequence selected from the group consisting of those sequences set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 5, 5b, 6b, 7, 8b, 9, 9b, 10, 10b, 11, 11b or 12.
- 20
12. A gene construct comprising a nucleotide sequence according to claims 7 to 11 operably linked to a heterologous nucleic acid.
13. A vector comprising a nucleotide sequence according to claims 7 to 11 or a
- 25
- gene construct according to claim 12.
14. A vector according to claim 13, which is an expression vector.
15. A vector according to claims 13 or 14, further comprising a selectable
- 30
- marker gene and/or nucleotide sequences for the replication in a host cell or the integration into the genome of the host cell.

16. A transgenic cell or organism containing a nucleotide sequence according to claims 7 to 11 and/or a gene construct according to claim 12 and/or a vector according to claims 13 to 15.
- 5
17. A transgenic cell or organism according to claim 16 which is an eucaryotic cell or organism.
18. A transgenic cell or organism according to claims 16 or 17 which is a yeast cell or a plant cell or a plant.
- 10
19. A transgenic cell or organism according to claims 16 to 18 having an altered biosynthetic pathway for the production of triacylglycerol.
- 15
20. A transgenic cell or organism according to claims 16 to 19 having an altered oil content.
21. A transgenic cell or organism according to claims 16 to 20 wherein the activity of PDAT is altered.
- 20
22. A transgenic cell or organism according to claims 16 to 21 wherein the altered activity of PDAT is characterized by an alteration in gene expression, catalytic activity and/or regulation of activity of the enzyme.
- 25
23. A transgenic cell or organism according to claims 16 to 22 wherein the altered biosynthetic pathway for the production of triacylglycerol is characterized by the prevention of accumulation of undesirable fatty acids in the membrane lipids.
- 30
24. A process for the production of triacylglycerol, comprising growing a transgenic cell or organism according to claims 16 to 23 under conditions

5

10

15

27. Use of a nucleotide sequence according to claims 7 to 11 and/or an enzyme according to claims 1 to 6 for the transformation of any cell or organism in order to be expressed in this cell or organism and result in an altered, preferably increased oil content of this cell or organism.

Abstract of the Disclosure

10

References cited in the description:

1. Bell, R. M. & Coleman, R. A. (1980) *Annu. Rev. Biochem.* **49**, 459-487.
2. Stymne, S. & Stobart, K. (1987) in *The biochemistry of plants: a comprehensive treatise*, Vol. 9, ed. Stumpf, P. K. (Academic Press, New York), pp. 175-214.
3. Cases, S. et al. (1998) *Proc. Natl. Acad. Sci. U S A* **95**, 13018-13023.
4. Hobbs, D. H., Lu, C. & Hills, M. J. (1999) *FEBS Lett.* **452**, 145-9
5. Zou, J., Wei, Y., Jako, C., Kumar, A., Selvaraj, G. & Taylor, D. C. (1999) *Plant J.* **19**, 645-653.
6. Lardizabal, K., Hawkins, D., Mai, J., & Wagner, N. (1999) Abstract presented at the Biochem. Mol. Plant Fatty Acids Glycerolipids Symposium, South Lake Tahoe, USA.
7. Thomas, B. J. & Rothstein, R. (1989) *Cell* **56**, 619-630.
8. Entian, K.-D. & Kötter, P. (1998) *Meth. Microbiol.* **26**, 431-449.
9. Kern, L., de Montigny, J., Jund, R. & Lacroute, F. (1990) *Gene* **88**, 149-157.
10. Ronne, H., Carlberg, M., Hu, G.-Z. & Nehlin, J. O. (1991) *Mol. Cell. Biol.* **11**, 4876-4884.
11. Stobart, K. & Stymne, S. (1990) in *Method in Plant Biochemistry*, vol 4, eds. Harwood, J. L. & Bowyer, J. R. (Academic press, London), pp. 19-46.
12. Bafor, M., Smith, M. A., Jonsson, L., Stobrt, A. K. & Stymne, S. (1991) *Biochem. J.* **280**, 507-514.
13. Banas, A., Johansson, I. & Stymne, S. (1992) *Plant Science* **84**, 137-144.
14. Kanda, P. & Wells, M. A. (1981) *J. Lipid. Res.* **22**, 877-879.
15. Ståhl, U., Ek, B. & Stymne, S. (1998) *Plant Physiol.* **117**, 197-205.
16. Stobart, K., Mancha, M. & Lenman M, Dahlqvist, A. & Stymne, S. (1997) *Planta* **203**, 58-66.
17. Bligh, E. G. & Dyer, W. J. (1959) *Can. J. Biochem. Physiol.* **37**, 911-917.
18. Sherman, F., Fink, G. R. & Hicks, J. B. (1986) in *Laboratory Course Manual for Methods in Yeast Genetics* (Cold Spring Harbor Laboratory)
19. Meesters, P. A. E. P., Huijberts, G. N. M. and Eggink, G. (1996) *Appl. Microbiol. Biotechnol.* **45**, 575-579.
20. van de Loo, F. J., Fox, B. G. & Sommerville, C. (1993), in *Lipid metabolism in plants*, ed. Moore, T. S. (CRC Press, Inc.), pp. 91-126.
21. van de Loo, F. J., Broun, P., Turner, S. & Sommerville, S. (1995) *Proc. Natl.*

Acad. Sci. U S A **95**, 6743-6747.

22. Lee, M., Lenman, M., Banas, A., Babor, M., Singh, S., Schweizer, M., Nilsson, R., Liljenberg, C., Dahlqvist, A., Gummesson, P-O., Sjö Dahl, S., Green, A., and Stymne, S. (1998) *Science* **280**, 915-918.
- 5 23. Glomset, J. A. (1968) *J. Lipid Res.* **9**, 155-167.
24. Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997) *Nucl. Acids Res.* **24**, 4876-4882.
25. Saitou, N. & Nei, M. (1987) *Mol. Biol. Evol.* **4**, 406-425.
26. Stålberg, K., Ellerström, M., Josefsson, L., & Rask, L. (1993) *Plant Mol. Biol.* **23**, 671
- 10 27. Becker, D., Kemper, E., Schell, J., Masterson, R. (1992) *Plant Mol. Biol.* **20**, 1195
28. D. Valvekens, M. Van Montagu, and Van Lusbettens (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 5536
- 15

Description of Figures

FIG. 1.

Metabolism of ^{14}C -labeled PC into the neutral lipid fraction by plant

5 microsomes. (A) Microsomes from developing seeds of sunflower, *R. communis* and *C. palaestina* were incubated for 80 min at 30°C with PC (8 nmol) having oleic acid in its *sn*-1 position, and either ^{14}C -labeled oleic, ricinoleic or vernolic acid in its *sn*-2 position. Radioactivity incorporated in TAG (open bars), DAG (solid bars), and unsterified fatty acids (hatched bars) was

10 quantified using thin layer chromatography followed by electronic autoradiography, and is shown as percentage of added labeled substrate. (B) Synthesis *in vitro* of TAG carrying two vernoloyl and one [^{14}C]ricinoleoyl group by microsomes from *R. communis*. The substrates added were unlabeled divernoloyl-DAG (5 nmol), together with either *sn*-1-oleoyl-*sn*-2-[^{14}C]ricinoleoyl-

15 DAG (0.4 nmol, 7700 dpm/nmol) or *sn*-1-oleoyl-*sn*-2-[^{14}C]ricinoleoyl-PC (0.4 nmol, 7700 dpm/nmol). The microsomes were incubated with the substrates for 30 min at 30°C, after which samples were removed for lipid analysis as described in the section „general methods“. The data shown are the average of two experiments.

20

FIG. 2.

PDAT activity in yeast microsomes, as visualized by autoradiogram of neutral lipid products separated on TLC.

Microsomal membranes (10 nmol of PC) from the wild type yeast strain FY1679 (lanes 1-3), a congeneric yeast strain (FVKT004-04C(AL)) that is disrupted for YNR008w (lane 4) or the same

25 disruption strain transformed with the plasmid pUS1, containing the YNR008w gene behind its native promotor (lane 5), were assayed for PDAT activity. As substrates, we used 2 nmol *sn*-1-oleoyl-*sn*-2-[^{14}C]ricinoleoyl-PC together with either 5 nmol of dioleoyl-DAG (lanes 2, 4 and 5) or *rac*-oleoyl-vernoleoyl-DAG

30 (lane 3). The enzymatic assay and lipid analysis was performed as described in Materials and Methods. The cells were precultured for 20 h in liquid YPD

medium, harvested and re-suspended in an equal volume of minimal medium (19) containing 16 g/l glycerol. The cells were then grown for an additional 24 h prior to being harvested. Selection for the plasmid was maintained by growing the transformed cells in synthetic medium lacking uracil (18). Abbreviations: 1-OH-TAG, monoricinoleoyl-TAG; 1-OH-1-ep-TAG, monoricinoleoyl-monovernoloyl-TAG; OH-FA, unesterified ricinoleic acid.

Fig. 3.

Lipid content (A,B) and PDAT activity (C) in PDAT overexpressing yeast cells.

10 The PDAT gene in the plasmid pUS4 was overexpressed from the galactose-induced *GAL1-TPK2* promotor in the wild type strain W303-1A (7). Its expression was induced after (A) 2 hours or (B) 25 hours of growth by the addition of 2% final concentration (w/v) of galactose. The cells were then incubated for another 22 hours before being harvested. The amount of lipids of

15 the harvested cells was determined by GLC-analysis of its fatty acid contents and is presented as μmol fatty acids per mg dry weight in either TAG (open bar), polar lipids (hatched bar), sterol esters (solid bar) and other lipids (striped bar). The data shown are the mean values of results with three independent yeast cultures. (C) *In vitro* synthesis of TAG by microsomes prepared from

20 yeast cells containing either the empty vector (vector) or the PDAT plasmid (+ PDAT). The cells were grown as in Fig. 3A. The substrate lipids dioleoyl-DAG (2.5 nmol) and *sn*-1-oleoyl-*sn*-2- ^{14}C -oleoyl-PC (2 nmol) were added to aliquots of microsomes (10 nmol PC), which were then incubated for 10 min at 28 °C. The amount of label incorporated into TAG was quantified by electronic

25 autoradiography. The results shown are the mean values of two experiments.

FIG. 4.

Substrate specificity of yeast PDAT. The PDAT activity was assayed by incubating aliquots of lyophilized microsomes (10 nmol PC) with substrate lipids at 30°C for 10 min (panel A) or 90 min (panel B). Unlabeled DAG (2.5 nmol)

30 was used as substrates together with different labeled phospholipids, as shown

in the figure. (A) *Sn*-position specificity of yeast PDAT regarding the acyl donor substrate. Dioleoyl-DAG together with either *sn*-1- ^{14}C oleoyl-*sn*-2- ^{14}C oleoyl-PC (di- ^{14}C -PC), *sn*-1- ^{14}C oleoyl-*sn*-2-oleoyl-PC (*sn*1- ^{14}C -PC) or *sn*-1-oleoyl-*sn*-2- ^{14}C oleoyl-PC (*sn*2- ^{14}C -PC). (B) Specificity of yeast PDAT regarding phospholipid headgroup and of the acyl composition of the phospholipid as well as of the diacylglycerol. Dioleoyl-DAG together with either *sn*-1-oleoyl-*sn*-2- ^{14}C oleoyl-PC (oleoyl-PC), *sn*-1-oleoyl-*sn*-2- ^{14}C oleoyl-PE (oleoyl-PE), *sn*-1-oleoyl-*sn*-2- ^{14}C ricinoleoyl-PC (ricinoleoyl-PC) or *sn*-1-oleoyl-*sn*-2- ^{14}C vernoleoyl-PC (vernoleoyl-PC). In the experiments presented in the 2 bars to the far right, monoricinoleoyl-DAG (ricinoleoyl-DAG or mono-vernoleoyl-DAG (vernoleoyl-DAG) were used together with *sn*-1-oleoyl-*sn*-2- ^{14}C oleoyl-PC. The label that was incorporated into TAG (solid bars) and lyso-PC (LPC, open bars) was quantified by electronic autoradiography. The results shown are the mean values of two experiments. The microsomes used were from W303-1A cells overexpressing the PDAT gene from the *GAL1-TPK2* promotor, as described in Fig. 3. The expression was induced at early stationary phase and the cells were harvested after an additional 24 h.

20 TAB.1:

In vitro synthesis of triacylglycerols in microsomal preparations of developing castor bean. Aliquots of microsomes (20 nmol PC) were lyophilised and substrate lipids were added in benzene solution: (A) 0.4 nmol ^{14}C -DAG (7760 dpm/nmol) and where indicated 1.6 nmol unlabelled DAG; (B) 0.4 nmol ^{14}C -DAG (7760 dpm/nmol) and 5 nmol unlabelled di-ricinoleoyl-PC and (C) 0.25 nmol ^{14}C -PC (4000 dpm/nmol) and 5 nmol unlabelled DAG. The benzene was evaporated by N_2 and 0.1 ml of 50 mM potassium phosphate was added, thoroughly mixed and incubated at 30 °C for (A) 20 min.; (B) and (C) 30 min.. Assays were terminated by extraction of the lipids in chloroform. The lipids were then separated by thin layer chromatography on silica gel 60 plates

(Merck; Darmstadt, Germany) in hexan/diethylether/acetic 35:70:1.5. The radioactive lipids were visualised and the radioactivity quantified on the plate by electronic autoradiography (Instant Imager, Packard, US). Results are presented as mean values of two experiments.

5

Radioactivity in different triacylglycerols (TAG) species formed. Abbreviations used: 1-OH-, mono-ricinoleoyl-; 2-OH-, di-ricinoleoyl-; 3-OH-, triricinoleoyl-; 1-OH-1-ver-, mono-ricinoleoyl-monovernoleoyl-; 1-OH-2-ver-, mono-ricinoleoyl-divernoleoyl-. Radiolabelled DAG and PC were prepared enzymatically. The radiolabelled ricinoleoyl group is attached at the sn-2-position of the lipid and unlabelled oleoyl group at the sn-1-position. Unlabelled DAG with vernoleoyl- or ricinoleoyl chains were prepared by the action of TAG lipase (6) on oil of *Euphorbia lagascae* or Castor bean, respectively. Synthetic di-ricinoleoyl-PC was kindly provided from Metapontum Agribios (Italy).

15

TAB.2:

Total fatty acids per mg of T2 seeds pooled from individual *Arabidopsis thaliana* plants transformed with yeast PDAT gene under the control of napin promotor (26-14) or transformed with empty vector (32-4).

20

* = statistical difference between control plants and PDAT transformed plants in a mean difference two-sided test at $\alpha = 5$.

Description of the SEQ ID:

5

SEQ ID NO. 2: The amino acid sequence of the suggested open reading frame YNR008w from *Saccharomyces cerevisiae*.

10

SEQ ID NO. 4: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with GenBank accession number AB006704.

SEQ ID NO. 5: Nucleotide sequence of the *Arabidopsis thaliana* cDNA clone with GenBank accession number T04806, and nucleotide ID number 315966.

20

SEQ ID NO. 7: Nucleotide and amino acid sequence of the *Zea mays* EST clone with GenBank accession number AI491339, and nucleotide ID number g4388167.

25

30

SEQ ID NO. 11: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with GenBank accession number AC003027.

10 SEQ ID NO. 13: Amino acid sequence of the *Schizosaccharomyces pombe*
putative open reading frame CAA22887 of the *Schizosaccharomyces pombe* gene
SPBC776.14.

15 *SEQ ID NO. 14*: Amino acid sequence of the *Arabidopsis thaliana* putative open reading frame AAC80628 derived from the *Arabidopsis thaliana* locus with GenBank accession number AC004557.

20 *SEQ ID NO 15*: Amino acid sequence of the *Arabidopsis thaliana* putative open reading frame AAD10668 derived from the *Arabidopsis thaliana* locus with GenBank accession number AC003027.


Further provisional and/or partial sequences are defined through the following SEQ IDs:

25 SEQ ID NO. 1a: The amino acid sequence of the yeast ORF YNR008w from *Saccharomyces cerevisiae*.

SEQ ID NO. 2a: Amino acid sequence of the region of the *Arabidopsis thaliana* genomic sequence (AC004557).

Figurs

Fig. 1:







Fig. 3:

Fig.4:



Tables

Tab. 1:

Substrate added	$[^{14}\text{C}]$ -lipid ⁽²⁾	unlabelled lipid ⁽²⁾	1-OH-TAG	mol % of added $[^{14}\text{C}]$ -acyl group in TAG ⁽¹⁾			
				2-OH-TAG	1-OH-1-ver-TAG	1-OH-2-ver-TAG	3-OH-TAG
A mono- $[^{14}\text{C}]$ -ricinoleoyl-DAG	mono-ricinoleoyl-DAG	mono-ricinoleoyl-DAG	2,8	12,4	-	-	-
A mono- $[^{14}\text{C}]$ -ricinoleoyl-DAG	mono-vernoleoyl-DAG	mono-vernoleoyl-DAG	3,2	12,1	1,3	-	-
A mono- $[^{14}\text{C}]$ -ricinoleoyl-DAG	di-vernoleoyl-DAG	di-vernoleoyl-DAG	4	10	0,5	1,2	-
A mono- $[^{14}\text{C}]$ -ricinoleoyl-DAG	di-ricinoleoyl-PC	di-ricinoleoyl-PC	0,3	24,8	-	-	-
B mono- $[^{14}\text{C}]$ -ricinoleoyl-PC	none	none	6,8	8,0	-	-	4,7
C mono- $[^{14}\text{C}]$ -ricinoleoyl-PC	di-oleoyl-DAG	di-oleoyl-DAG	8,6	9,8	-	-	5,0
C mono- $[^{14}\text{C}]$ -ricinoleoyl-PC	mono-ricinoleoyl-DAG	mono-ricinoleoyl-DAG	5,7	16,7	-	-	1,9
C mono- $[^{14}\text{C}]$ -ricinoleoyl-PC	di-ricinoleoyl-DAG	di-ricinoleoyl-DAG	4,5	9,4	-	-	9,5
C mono- $[^{14}\text{C}]$ -ricinoleoyl-PC	mono-vernoleoyl-DAG	mono-vernoleoyl-DAG	6,0	11,5	10,9	0,5	7,4
C mono- $[^{14}\text{C}]$ -ricinoleoyl-PC	di-vernoleoyl-DAG	di-vernoleoyl-DAG	6,7	10,8	1,1	8,4	6,8

Sequence Listing

<210> 1

<211> 1986

<212> genomic DNA

<213> *Saccharomyces cerevisiae*

<221> CDS

<222> (1)..(1983)

<400> 1

```

atg ggc aca ctg ttt cga aga aat gtc cag aac caa aag agt gat tct      48
Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
   1             5             10             15

gat gaa aac aat aaa ggg ggt tct gtt cat aac aag cga gag agc aga      96
Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
          20             25             30

aac cac att cat cat caa cag gga tta ggc cat aag aga aga agg ggt      144
Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
          35             40             45

att agt ggc agt gca aaa aga aat gag cgt ggc aaa gat ttc gac agg      192
Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
          50             55             60

aaa aga gac ggg aac ggt aga aaa cgt tgg aga gat tcc aga aga ctg      240
Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
          65             70             75             80

att ttc att ctt ggt gca ttc tta ggt gta ctt ttg ccg ttt agc ttt      288
Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
          85             90             95

ggc gct tat cat gtt cat aat agc gat agc gac ttg ttt gac aac ttt      336
Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
          100             105             110

gta aat ttt gat tca ctt aaa gtg tat ttg gat gat tgg aaa gat gtt      384
Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
          115             120             125

ctc cca caa ggt ata agt tcg ttt att gat gat att cag gct ggt aac      432
Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
          130             135             140

```

00000000000000000000

<210> 4
 <211> 3685
 <212> genomic DNA
 <213> Arabidopsis thaliana

<400> 4

ATGCCCTTA	TTCATCGGAA	AAAGCCGACG	GAGAAACCAT	CGACGCCGCC	50
ATCTGAAGAG	GTGGTGCACG	ATGAGGATTC	GCAAAAGAAA	CCACACGAAT	100
CTTCCAAATC	CCACCATAAG	AAATCGAACG	GAGGAGGGAA	GTGGTCGTGC	150
ATCGATTCTT	GTGTGTGGTT	CATTGGGTGT	GTGTGTGTAA	CCTGGTGGTT	200
TCTTCTCTTC	CTTTACAACG	CAATGCCTGC	GAGCTTCCCT	CAGTATGTAA	250
CGGAGCGAAT	CACGGGTCTT	TTGCCTGACC	CGCCCGGTGT	TAAGCTCAAA	300
AAAGAAGGTC	TAAAGCGGAA	ACATCCTGTT	GTCTTCATTC	CTGGGATTGT	350
CACCGGTGGG	CTCGAGCTTT	GGGAAGGCAA	ACAATGCGCT	GATGGTTTAT	400
TTAGAAAACG	TTTGTGGGGT	GGAACCTTTG	GTGAAGTCTA	CAAAAGGTGA	450
GCTCAACAAT	TCTCACTCTT	CCTTTATATT	GGGATTTGGA	TTGGATCTGA	500
TGAGATCACG	CACCTGTGTC	TTCTTCAACA	TCACTCAAAC	TTTAATTCCA	550
TGTTTGTCTG	TCTTACTCTT	TACTTTTTTT	TTTTTTTGAT	GTGAAACGCT	600
ATTTTCTTAA	GAGACTATTT	CTGTATGTGT	AAGGTAAGCG	TTCCAAGGAC	650
GTAATTGGCT	TGGACTATTT	CTGTTTGATT	GTAACTTTTA	GGATATAAAA	700
TAGCTGCCTT	GGAATTTCAA	GTCATCTTAT	TGCCAAATCT	GTTGCTAGAC	750
ATGCCCTAGA	GTCCGTTCAT	AACAAGTTAC	TTCCCTTACT	GTCGTTGCGT	800
GTAGATTTAG	CTTTGTGTAG	CGTATAATGA	AGTAGTGTTC	TATGTTTGTG	850
TGGGAATAGA	GAAGTTCTAA	CTACATCTGT	GGAAAGTGTG	TTCAGGCTGT	900
GATAGAGGAC	TGTTGCTTTA	TTATTCAACT	ATGTATATGT	GTAATTAAAG	950
CTAGTTCCCT	TTTGATCTTT	CAGCTCAATG	TGCTTTTCTC	AATTTTTTTC	1000
TCAATTTCAA	AGTTTCACAT	CGAGTTTATT	CACATGTCTT	GAATTTTCGT	1050
CATCCTCGTT	CTGTTATCCA	GCTTTGAACT	CCTCCCGACC	CTGCTATGGA	1100
TATATTAAAA	AAAAAGTGTG	TTGTGGGTTG	CATCTTTGTT	ACGATCTGCA	1150
TCTTCTCTCT	TCCGCTCAGT	GTTTATGTTT	TTGCTATGGT	AGAGATGGGC	1200
AATGTTATTG	TTGATGGTAA	CAGTGGTATA	GTTGATAGTA	TCTTAACTAA	1250
TCAATTATCT	CTTTGATTCA	GGCCTCTATG	TTGGGTGGAA	CACATGTCAC	1300
TTGACAATGA	AACTGGTTTG	GATCCAGCTG	GTATTAGAGT	TCCAGCTGTA	1350
TCAGGACTCG	TGGCTGCTGA	CTACTTTGCT	CCTGGCTACT	TTGTCTGGGC	1400
AGTGCTGATT	GCTAACCTTG	CACATATTGG	ATATGAAGAG	AAAAATATGT	1450
ACATGGCTGC	ATATGACTGG	CGGCTTTCGT	TTCAGAACAC	AGAGGTTCTT	1500
TTCTCATCGT	TCTTTCTATT	ATTCTGTTCC	ATGTTACGTT	TCTTTCTTCA	1550
TTACTTAAGG	CTTAAATATG	TTTCATGTTG	AATTAATAGG	TACGTGATCA	1600
GACTCTTAGC	CGTATGAAAA	GTAATATAGA	GTTGATGGTT	TCTACCAACG	1650
GTGGAATAAA	AGCAGTTATA	GTTCCGCATG	CCATGGGGGT	CTTGATTTTT	1700
CTACATTTTA	TGAAGTGGGT	TGAGGCACCA	GCTCCTCTGG	GTGGCGGGGG	1750
TGGGCCAGAT	TGGTGTGCAA	AGTATATTAA	GGCGGTGATG	AACATTGGTG	1800
GACCATTCTT	TGGTGTTCCT	AAAGCTGTTG	CAGGGCTTTT	CTCTGCTGAA	1850
GCAAAGGATG	TTGCAGTTGC	CAGGTATTGA	ATATCTGCTT	ATACTTTTGA	1900
TGATCAGAAC	CTTGGCTCTG	GAACTCAAAG	TTATTCTACT	AAATATCAAT	1950
TCTAATAACA	TTGCTATATT	ATCGCTGCAA	CTGACATTGG	TTGATTATTT	2000
TTGCTGCTTA	TGTAAGTAA	ACTCTCTTGA	GATTAGACAA	ATGATGAATT	2050
GATAATTCTT	ACGCATTGCT	CTGTGATGAC	CAGTTTCTTA	GCTTCGACGA	2100
TAACATTTGT	CATACTGTCT	TTTGGAGGGC	ATTGAATTTT	GCTATGGAAA	2150
GCGCTGGAGC	TTCCATGCTT	GCATTCTTTA	CCAATTAGCG	TTATTCTGCT	2200
TCTTTCAATT	TTCTTGTATA	TGCATCTATG	GTCTTTTATT	TCTTCTTAAT	2250
TAAAGACTCG	TTGGATTAGT	TGCTCTATTA	GTCACCTGGT	TCCTTAATAT	2300
AGAACTTTAC	TTTCTTCGAA	AATTGCAGAG	CGATTGCCCC	AGGATTCTTA	2350
GACACCGATA	TATTTAGACT	TCAGACCTTG	CAGCATGTAA	TGAGAATGAC	2400
ACGCACATGG	GACTCAACAA	TGTCTATGTT	ACCGAAGGGA	GGTGACACGA	2450
TATGGGGCGG	GCTTGATTGG	TCACCGGAGA	AAGGCCACAC	CTGTTGTGGG	2500
AAAAAGCAAA	AGAACAACGA	AACTTGTGGT	GAAAGCAGGTG	AAAACGGAGT	2550
TTCCAAGAAA	AGTCTGTGTA	ACTATGGAAG	GATGATATCT	TTTGGGAAAG	2600
AAGTAGCAGA	GGCTGCGCCA	TCTGAGATTA	ATAATATTGA	TTTTTCGAGTA	2650
AGGACATATA	AATCATAATA	AACCTTGTAC	ATTTTGTGAT	TGTATGATGA	2700
ATATCTGTAC	ATTTTATCTG	GTGAAGGGTG	CTGTCAAAGG	TCAGAGTATC	2750
CCAAATCACA	CCTGTCTGTA	CGTGTGGACA	GAGTACCATG	ACATGGGAAT	2800
TGCTGGGATC	AAAGCTATCG	CTGAGTATAA	GGTCTACACT	GCTGGTGAAG	2850

00000-0122550

<210> 7
 <211> 643
 <212> cDNA
 <213> Zea mays

<221> CDS
 <222> (1)..(402)

<400> 6
 CGG GAG AAA ATA GCT GCT TTG AAG GGG GGT GTT TAC TTA GCC GAT GGT 48
 Arg Glu Lys Ile Ala Ala Leu Lys Gly Gly Val Tyr Leu Ala Asp Gly
 1 5 10 15

GAT GAA ACT GTT CCA GTT CTT AGT GCG GGC TAC ATG TGT GCG AAA GGA 96
 Asp Glu Thr Val Pro Val Leu Ser Ala Gly Tyr Met Cys Ala Lys Gly
 20 25 30

TGG CGT GGC AAA ACT CGT TTC AGC CCT GCC GGC AGC AAG ACT TAC GTG 144
 Trp Arg Gly Lys Thr Arg Phe Ser Pro Ala Gly Ser Lys Thr Tyr Val
 35 40 45

AGA GAA TAC AGC CAT TCG CCA CCC TCT ACT CTC CTG GAA GGC AGG GGC 192
 Arg Glu Tyr Ser His Ser Pro Pro Ser Thr Leu Leu Glu Gly Arg Gly
 50 55 60

ACC CAG AGC GGT GCA CAT GTT GAT ATA ATG GGG AAC TTT GCT CTA ATT 240
 Thr Gln Ser Gly Ala His Val Asp Ile Met Gly Asn Phe Ala Leu Ile
 65 70 75 80

GAG GAC GTC ATC AGA ATA GCT GCT GGG GCA ACC GGT GAG GAA ATT GGT 288
 Glu Asp Val Ile Arg Ile Ala Ala Gly Ala Thr Gly Glu Glu Ile Gly
 85 90 95

GGC GAT CAG GTT TAT TCA GAT ATA TTC AAG TGG TCA GAG AAA ATC AAA 336
 Gly Asp Gln Val Tyr Ser Asp Ile Phe Lys Trp Ser Glu Lys Ile Lys
 100 105 110

TTG AAA TTG TAA CCTATGGGAA GTTAAAGAAG TGCCGACCCG TTTATTGCGTTCC 391
 Leu Lys Leu
 115

AAAGTGTCTT GCCTGAGTGC AACTCTGGAT TTTGCTTAAA TATTGTAATT TTTCACGC 449
 TTCATTCTGTC CCTTTGTCAA ATTTACATTT GACAGGACGC CAATGCGATA CGATGTTG 507
 TACCGCTATT TTCAGCATTG TATATTAAAC TGTACAGGTG TAAGTTGCAT TTGCCAGC 565
 TGAAATTGTG TAGTCGTTTT CTTTACGATT TAATANCAAG TGGCGGAGCA GTGCCCCA 623
 AGCNAAAAAA AAAAAAAAAA 643

CGG GAG AAA ATA GCT GCT TTG AAG GGG GGT GTT TAC TTA GCC GAT GGT 48

3896

<400> 12

CTGGGGCCAA	AAGTGAACAT	AACAAGGACA	CCACAGTCAG	AGCATGATGT	50
TCAGATGTAC	AAGTGCATCT	AAATATAGAG	CATCAACATG	GTGAAGATAT	100
CATTCCCAAT	ATGACAAAGT	TACCTACAAT	GAAGTACATA	ACCTATTATG	150
AGGATTCTGA	AAGTTTTCCA	GGGACAAGAA	CAGCAGTTTG	GGAGCTTGAT	200
AAAGCAAATC	ACAGGAACAT	TGTCAGATCT	CCAGCTTTGA	TGCGGGAGCT	250
TGGGCTTGAG	ATGTGGCATG	ATATTTCATC	TGATAAAATG	TCCAAGTTTG	300
TTACAAAAGG	TGGTGTCTGA	TCCTCACTAT	TTTCTTCTAT	AAATGTTTGA	350
GTTTGTATTG	ACATTGTAAG	TATTGCAACA	AAAAGCAAAG	CGTGGGCCTC	400
TGAGGGATGA	GGACTGCTAT	TGGGATTACG	GGAAAGCTCG	ATGTGCATGG	450
GCTGAACATT	GTGAATACAG	GTTAGAATAT	TCAAATTATA	TTTTGCAAAA	500
TATTCTCTTT	TTGTGTATTT	AGGCCACCTT	TCCCCGGTCA	CAACGATGCA	550
GATATGTATT	CGGGGATGTT	CACCTGGGAC	AGAGTTGCAG	ATTGAAGAGT	600
TCTACATCTC	ACATCCTGTC	ACACTATGTG	TGATATTTAA	GAAACTTTGT	650
TTGGCGGAAC	AACAAGTTTG	CACAAACATT	TGAAGAAGAA	AGCGAAATGA	700
TTCAGAGAG					

709

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

<400> 13

MASSKSKTHKKKEVKSPIDLNSKKPTRLALSEQPSASETQSVSNKSRKSKFGKRLNFI LGAILGICGA 70
FFFAVGDDNAVFDPATLDKFGNMLGSSDLFDDIKGYLSYNVFKDAPFTTDKPSQSPSGNEVQVGLDMYNE 140
GYRSDHPVIMVPGVISSGLESWSFNNCSI PYFRKRLWGSWSMLKAMFLDKQCWLEHMLDKKTGLDPKGI 210
KLRAAQGFEEADFFITGYWIWSKVIENTLAAIGYEPNNMLSASYDWRLSYANLEERDKYFSKLKMFIEYSN 280
IVHKKKVVLISHSMGSQVTTYFFKWVEAEGYGNGGPTWVNDHIEAFINISGSLIGAPKTVAAALLSGEMKD 350
TGIVITLNLILEKFFSRSERAMMVRTMGGVSSMLPKGGDVAPDDL NQTNFSNGAIIRYREDIDKDHDEFDI 420
DDALQFLKNVTDDDFKVMLAKNYSHGLAWTEKEVLKNNEMPSKWINPLETSLPYAPDMKIYCVHGVGKPT 490
ERGYYYTNNPEGQFVIDSSVNDGTKVENGIVMDDGDGTLPIALGLVCNKVWQTKRFNPANTSITNYEIK 560
HEPAADFRLRGGRSAEHVDILGHSELNEIILKVSSGHGDSVPNRYISDIOEIIINEINLDKPRN 623

Parameter	Value	Unit
α_1	0.1	
α_2	0.1	
α_3	0.1	
α_4	0.1	
α_5	0.1	
α_6	0.1	
α_7	0.1	
α_8	0.1	
α_9	0.1	
α_{10}	0.1	
α_{11}	0.1	
α_{12}	0.1	
α_{13}	0.1	
α_{14}	0.1	
α_{15}	0.1	
α_{16}	0.1	
α_{17}	0.1	
α_{18}	0.1	
α_{19}	0.1	
α_{20}	0.1	
α_{21}	0.1	
α_{22}	0.1	
α_{23}	0.1	
α_{24}	0.1	
α_{25}	0.1	
α_{26}	0.1	
α_{27}	0.1	
α_{28}	0.1	
α_{29}	0.1	
α_{30}	0.1	
α_{31}	0.1	
α_{32}	0.1	
α_{33}	0.1	
α_{34}	0.1	
α_{35}	0.1	
α_{36}	0.1	
α_{37}	0.1	
α_{38}	0.1	
α_{39}	0.1	
α_{40}	0.1	
α_{41}	0.1	
α_{42}	0.1	
α_{43}	0.1	
α_{44}	0.1	
α_{45}	0.1	
α_{46}	0.1	
α_{47}	0.1	
α_{48}	0.1	
α_{49}	0.1	
α_{50}	0.1	
α_{51}	0.1	
α_{52}	0.1	
α_{53}	0.1	
α_{54}	0.1	
α_{55}	0.1	
α_{56}	0.1	
α_{57}	0.1	
α_{58}	0.1	
α_{59}	0.1	
α_{60}	0.1	
α_{61}	0.1	
α_{62}	0.1	
α_{63}	0.1	
α_{64}	0.1	
α_{65}	0.1	
α_{66}	0.1	
α_{67}	0.1	
α_{68}	0.1	
α_{69}	0.1	
α_{70}	0.1	
α_{71}	0.1	
α_{72}	0.1	
α_{73}	0.1	
α_{74}	0.1	
α_{75}	0.1	
α_{76}	0.1	
α_{77}	0.1	
α_{78}	0.1	
α_{79}	0.1	
α_{80}	0.1	
α_{81}	0.1	
α_{82}	0.1	
α_{83}	0.1	
α_{84}	0.1	
α_{85}	0.1	
α_{86}	0.1	
α_{87}	0.1	
α_{88}	0.1	
α_{89}	0.1	
α_{90}	0.1	
α_{91}	0.1	
α_{92}	0.1	
α_{93}	0.1	
α_{94}	0.1	
α_{95}	0.1	
α_{96}	0.1	
α_{97}	0.1	
α_{98}	0.1	
α_{99}	0.1	
α_{100}	0.1	

```
<210> 14
<211> 432
<212> PRT
<213> Arabidopsis thaliana
```

```
<400> 14
MKKISSHYSVVIAILVVVTMTSMCQAVGSNNVYPLILVPGNGGNQLEVRLDREYKPSVWCVSSWLYPIHKK 70
SGGWFRWFDAAVLLSPFTRCFSDRMMLYYDPDLDDYQNA PGVQTRVPHFGSTKSLLYLDPRLRDATSYM 140
EHLVKALEKKCGYVNDQITLGAPYDFRYGLAASGHPSRVASQFLQDLKQLVEKTSSENEKGPVILLSHSL 210
GGLFVLHFLNRTTPSWRRKYIKHFVALAAPWG GTISQMKTFASGNTLGVPLVNPLLVRRHQRTSESNQWL 280
LPSTKVFHDRTKPLVVTPQVNYTAYEMDRFFADIGFSQGVVPYKTRVLPLTEELMTPGVPVTCIYGRGVD 350
TPEVLMYGKG GFDKQPEIKYGDGDGTVN LASLAALKVDSLNTVEIDGVSHTSILKDEIALKEIMKQISII 420
NYELANVNAVNE
432
```

432

[illegible]

<210> 15

<211> 552

<212> PRT

<213> *Arabidopsis thaliana*

<400> 15

MGANSKSVTASFTVIAVFFLICGGRTAVEDETEFHGDYSKLSGIIIPGFASTQLRAWSILDCPYTPLDFN 70
PLDLVWLDTTKLLSAVNCWFKCMVLDPNQTDHPECKSRPDSGLSAITELDPGYITGPLSTVWKEWLKWC 140
VEFGIEANAIVAVPYDWRLSPTKLEERDLYFHKLKLTFETALKLRGGPSIVFAHSMGNNVFRYFLEWLRL 210
EIAPKHYLKWLDQHIHAYFAVGAPLLGSVEAIKSTLSGVTFGLPVSEGTARLLSNSFASSLWLMPFSKNC 280
KGDNTFWTHFSGGAACKDKRVYHCDEEEYQSKYSGWPTNIINIEIPSTSARELADGTLFKAIEDYDPSK 350
RMLHQLKKYVPFFVIRNIAHRSSLAGFLLYHDDPVFNPLTPWERPPIKNVFCIYGAHLKTEVGYYFAPSG 420
KPYPDNWIITDIIYETEGSLVSRSGTVVDGNAGPITGDETVPYHSLSWCKNWLGPKNITMAPQILIGKI 490
KQQPEHDGSDVHVELNVDHEHGSDIIANMTKAPRVKYITFYEDSESIPGKRTAVWELDKSGY 552

[illegible]

5

10

15

<170> PatentIn Ver. 2.0

<210> 1a

<211> 661

20 <212> PRT

<213> Saccharomyces cerevisiae

<400> 1

25 Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
1 5 10 15Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
20 25 3030 Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
35 40 45Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
50 55 6035 Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
65 70 75 8040 Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
85 90 95Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
100 105 11045 Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
115 120 125Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
130 135 14050 Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly
145 150 155 16055 Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val
165 170 175Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile
180 185 19060 Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp
195 200 205

"GZ666"

	Gly 210	Ser	Phe	Tyr	Met	Leu	Arg 215	Thr	Met	Val	Met 220	Asp	Lys	Val	Cys	Trp
5	Leu 225	Lys	His	Val	Met	Leu 230	Asp	Pro	Glu	Thr	Gly 235	Leu	Asp	Pro	Pro	Asn 240
10	Phe	Thr	Leu	Arg	Ala 245	Ala	Gln	Gly	Phe	Glu 250	Ser	Thr	Asp	Tyr	Phe 255	Ile
	Ala	Gly	Tyr	Trp 260	Ile	Trp	Asn	Lys	Val 265	Phe	Gln	Asn	Leu	Gly 270	Val	Ile
15	Gly	Tyr	Glu 275	Pro	Asn	Lys	Met 280	Thr	Ser	Ala	Ala	Tyr	Asp 285	Trp	Arg	Leu
	Ala	Tyr 290	Leu	Asp	Leu	Glu	Arg 295	Arg	Asp	Arg	Tyr	Phe 300	Thr	Lys	Leu	Lys
20	Glu 305	Gln	Ile	Glu	Leu	Phe 310	His	Gln	Leu	Ser	Gly 315	Glu	Lys	Val	Cys	Leu 320
25	Ile	Gly	His	Ser	Met 325	Gly	Ser	Gln	Ile	Ile 330	Phe	Tyr	Phe	Met	Lys 335	Trp
	Val	Glu	Ala 340	Glu	Gly	Pro	Leu	Tyr	Gly 345	Asn	Gly	Gly	Arg	Gly 350	Trp	Val
30	Asn	Glu	His 355	Ile	Asp	Ser	Phe 360	Ile	Asn	Ala	Ala	Gly	Thr 365	Leu	Leu	Gly
	Ala 370	Pro	Lys	Ala	Val	Pro	Ala 375	Leu	Ile	Ser	Gly	Glu 380	Met	Lys	Asp	Thr
35	Ile 385	Gln	Leu	Asn	Thr	Leu 390	Ala	Met	Tyr	Gly	Leu 395	Glu	Lys	Phe	Phe	Ser 400
40	Arg	Ile	Glu	Arg	Val 405	Lys	Met	Leu	Gln	Thr 410	Trp	Gly	Gly	Ile	Pro	Ser
	Met	Leu	Pro	Lys 420	Gly	Glu	Glu	Val	Ile 425	Trp	Gly	Asp	Met	Lys 430	Ser	Ser
45	Ser	Glu	Asp 435	Ala	Leu	Asn	Asn 440	Thr	Asp	Thr	Tyr	Gly 445	Asn	Phe	Ile	
	Arg 450	Phe	Glu	Arg	Asn	Thr	Ser 455	Asp	Ala	Phe	Asn	Lys 460	Asn	Leu	Thr	Met
50	Lys 465	Asp	Ala	Ile	Asn	Met 470	Thr	Leu	Ser	Ile	Ser 475	Pro	Glu	Trp	Leu	Gln 480
55	Arg	Arg	Val	His 485	Glu	Gln	Tyr	Ser	Phe	Gly 490	Tyr	Ser	Lys	Asn	Glu 495	Glu
	Glu	Leu	Arg	Lys 500	Asn	Glu	Leu	His	His 505	Lys	His	Trp	Ser	Asn 510	Pro	Met
60	Glu	Val	Pro 515	Leu	Pro	Glu	Ala 520	Pro	His	Met	Lys	Ile 525	Tyr	Cys	Ile	Tyr

	Gly	Val	Asn	Asn	Pro	Thr	Glu	Arg	Ala	Tyr	Val	Tyr	Lys	Glu	Glu	Asp
	530						535						540			
5	Asp	Ser	Ser	Ala	Leu	Asn	Leu	Thr	Ile	Asp	Tyr	Glu	Ser	Lys	Gln	Pro
	545					550					555					560
	Val	Phe	Leu	Thr	Glu	Gly	Asp	Gly	Thr	Val	Pro	Leu	Val	Ala	His	Ser
					565					570					575	
10	Met	Cys	His	Lys	Trp	Ala	Gln	Gly	Ala	Ser	Pro	Tyr	Asn	Pro	Ala	Gly
				580					585					590		
	Ile	Asn	Val	Thr	Ile	Val	Glu	Met	Lys	His	Gln	Pro	Asp	Arg	Phe	Asp
15			595					600					605			
	Ile	Arg	Gly	Gly	Ala	Lys	Ser	Ala	Glu	His	Val	Asp	Ile	Leu	Gly	Ser
	610						615					620				
20	Ala	Glu	Leu	Asn	Asp	Tyr	Ile	Leu	Lys	Ile	Ala	Ser	Gly	Asn	Gly	Asp
	625					630					635					640
	Leu	Val	Glu	Pro	Arg	Gln	Leu	Ser	Asn	Leu	Ser	Gln	Trp	Val	Ser	Gln
					645					650					655	
25	Met	Pro	Phe	Pro	Met											
				660												
30	<210> 2Q															
	<211> 387															
	<212> PRT															
	<213> Arabidopsis thaliana															
35	<400> 2															
	Val	Gly	Ser	Asn	Val	Tyr	Pro	Leu	Ile	Leu	Val	Pro	Gly	Asn	Gly	Gly
	1				5					10					15	
	Asn	Gln	Leu	Glu	Val	Arg	Leu	Asp	Arg	Glu	Tyr	Lys	Pro	Ser	Ser	Val
40				20					25					30		
	Trp	Cys	Ser	Ser	Trp	Leu	Tyr	Pro	Ile	His	Lys	Lys	Ser	Gly	Gly	Trp
		35					40						45			
45	Phe	Arg	Leu	Trp	Phe	Asp	Ala	Ala	Val	Leu	Leu	Ser	Pro	Phe	Thr	Arg
	50						55					60				
	Cys	Phe	Ser	Asp	Arg	Met	Met	Leu	Tyr	Tyr	Asp	Pro	Asp	Leu	Asp	Asp
	65					70					75					80
50	Tyr	Gln	Asn	Ala	Pro	Gly	Val	Gln	Thr	Arg	Val	Pro	His	Phe	Gly	Ser
					85					90					95	
	Thr	Lys	Ser	Leu	Leu	Tyr	Leu	Asp	Pro	Arg	Leu	Arg	Asp	Ala	Thr	Ser
55				100					105					110		
	Tyr	Met	Glu	His	Leu	Val	Lys	Ala	Leu	Glu	Lys	Lys	Cys	Gly	Tyr	Val
			115					120								

	165								170								175								
5	atg	ggt	cct	ggt	gtc	att	tct	acg	gga	att	gaa	agc	tgg	gga	gtt	att	576								
	Met	Val	Pro	Gly	Val	Ile	Ser	Thr	Gly	Ile	Glu	Ser	Trp	Gly	Val	Ile									
				180					185					190											
10	gga	gac	gat	gag	tgc	gat	agt	tct	gcg	cat	ttt	cgt	aaa	cgg	ctg	tgg	624								
	Gly	Asp	Asp	Glu	Cys	Asp	Ser	Ser	Ala	His	Phe	Arg	Lys	Arg	Leu	Trp									
			195					200					205												
15	gga	agt	ttt	tac	atg	ctg	aga	aca	atg	ggt	atg	gat	aaa	gtt	tgt	tgg	672								
	Gly	Ser	Phe	Tyr	Met	Leu	Arg	Thr	Met	Val	Met	Asp	Lys	Val	Cys	Trp									
		210					215					220													
20	ttg	aaa	cat	gta	atg	tta	gat	cct	gaa	aca	ggt	ctg	gac	cca	ccg	aac	720								
	Leu	Lys	His	Val	Met	Leu	Asp	Pro	Glu	Thr	Gly	Leu	Asp	Pro	Pro	Asn									
	225					230					235					240									
25	ttt	acg	cta	cgt	gca	gca	cag	ggc	ttc	gaa	tca	act	gat	tat	ttc	atc	768								
	Phe	Thr	Leu	Arg	Ala	Ala	Gln	Gly	Phe	Glu	Ser	Thr	Asp	Tyr	Phe	Ile									
					245				250						255										
30	gca	ggg	tat	tgg	att	tgg	aac	aaa	ggt	ttc	caa	aat	ctg	gga	gta	att	816								
	Ala	Gly	Tyr	Trp	Ile	Trp	Asn	Lys	Val	Phe	Gln	Asn	Leu	Gly	Val	Ile									
				260					265					270											
35	ggc	tat	gaa	ccc	aat	aaa	atg	acg	agt	gct	gcg	tat	gat	tgg	agg	ctt	864								
	Gly	Tyr	Glu	Pro	Asn	Lys	Met	Thr	Ser	Ala	Ala	Tyr	Asp	Trp	Arg	Leu									
			275					280					285												
40	gca	tat	tta	gat	cta	gaa	aga	cgc	gat	agg	tac	ttt	acg	aag	cta	aag	912								
	Ala	Tyr	Leu	Asp	Leu	Glu	Arg	Arg	Asp	Arg	Tyr	Phe	Thr	Lys	Leu	Lys									
		290					295					300													
45	gaa	caa	atc	gaa	ctg	ttt	cat	caa	ttg	agt	ggt	gaa	aaa	gtt	tgt	tta	960								
	Glu	Gln	Ile	Glu	Leu	Phe	His	Gln	Leu	Ser	Gly	Glu	Lys	Val	Cys	Leu									
	305					310					315					320									
50	att	gga	cat	tct	atg	ggt	tct	cag	att	atc	ttt	tac	ttt	atg	aaa	tgg	1008								
	Ile	Gly	His	Ser	Met	Gly	Ser	Gln	Ile	Ile	Phe	Tyr	Phe	Met	Lys	Trp									
					325				330						335										
55	gtc	gag	gct	gaa	ggc	cct	ctt	tac	ggt	aat	ggt	ggt	cgt	ggc	tgg	gtt	1056								
	Val	Glu	Ala	Glu	Gly	Pro	Leu	Tyr	Gly	Asn	Gly	Gly	Arg	Gly	Trp	Val									
				340					345					350											
60	aac	gaa	cac	ata	gat	tca	ttc	att	aat	gca	gca	ggg	acg	ctt	ctg	ggc	1104								
	Asn	Glu	His	Ile	Asp	Ser	Phe	Ile	Asn	Ala	Ala	Gly	Thr	Leu	Leu	Gly									
			355					360					365												
65	gct	cca	aag	gca	gtt	cca	gct	cta	att	agt	ggt	gaa	atg	aaa	gat	acc	1152								
	Ala	Pro	Lys	Ala	Val	Pro	Ala	Leu	Ile	Ser	Gly	Glu	Met	Lys	Asp	Thr									
		370					375					380													
70	att	caa	tta	aat	acg	tta	gcc	atg	tat	ggt	ttg	gaa	aag	ttc	ttc	tca	1200								
	Ile	Gln	Leu	Asn	Thr	Leu	Ala	Met	Tyr	Gly	Leu	Glu	Lys	Phe	Phe	Ser									
	385					390					395					400									
75	aga	att	gag	aga	gta	aaa	atg	tta	caa	acg	tgg	ggt	ggt	ata	cca	tca	1248								
	Arg	Ile	Glu	Arg	Val	Lys	Met	Leu	Gln	Thr	Trp	Gly	Gly	Ile	Pro	Ser									
					405					410					415										

Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp
 595 600 605
 5 Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser
 610 615 620
 Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp
 625 630 635 640
 10 Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln
 645 650 655
 Met Pro Phe Pro Met
 660
 15

666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000

gca tat tta gat cta gaa aga cgc gat agg tac ttt acg aag cta aag 912
 Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys
 290 295 300

gaa caa atc gaa ctg ttt cat caa ttg agt ggt gaa aaa gtt tgt tta 960
 Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu
 305 310 315 320

att gga cat tct atg ggt tct cag att atc ttt tac ttt atg aaa tgg 1008
 Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp
 325 330 335

gtc gag gct gaa ggc cct ctt tac ggt aat ggt ggt cgt ggc tgg gtt 1056
 Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val
 340 345 350

aac gaa cac ata gat tca ttc att aat gca gca ggg acg ctt ctg ggc 1104
 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly
 355 360 365

gct cca aag gca gtt cca gct cta att agt ggt gaa atg aaa gat acc 1152
 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr
 370 375 380

att caa tta aat acg tta gcc atg tat ggt ttg gaa aag ttc ttc tca 1200
 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser
 385 390 395 400

aga att gag agn gta aaa atg tta caa acg tgg ggt ggt ata cca tca 1248
 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser
 405 410 415

atg cta cca aag gga gaa gag gtc att tgg ggg gat atg aag tca tct 1296
 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser
 420 425 430

tca gag gat gca ttg aat aac aac act gac aca tac ggc aat ttc att 1344
 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile
 435 440 445

cga ttt gaa agg aat acg agc gat gct ttc aac aaa aat ttg aca atg 1392
 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met
 450 455 460

aaa gac gcc att aac atg aca tta tgg ata tca cct gaa tgg ctc caa 1440
 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln
 465 470 475 480

000000-000000

<210> 219

<211> 661

<212> PRT

<213> *Saccharomyces cerevisiae*

<400> 2

Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
 1 5 10 15
 Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
 20 25 30
 Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
 35 40 45
 Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
 50 55 60
 Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
 65 70 75 80
 Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
 85 90 95
 Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
 100 105 110
 Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
 115 120 125
 Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
 130 135 140
 Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly
 145 150 155 160
 Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val
 165 170 175
 Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile
 180 185 190
 Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp
 195 200 205
 Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp
 210 215 220
 Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn
 225 230 235 240
 Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile
 245 250 255
 Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile
 260 265 270
 Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu
 275 280 285
 Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys
 290 295 300
 Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu
 305 310 315 320

00000-01220000

<210> 5
 <211> 402
 <212> cDNA
 <213> Arabidopsis thaliana
 <220>
 <221> CDS
 <222> (120)..(402)
 <400> 5

AGAAACAGCTCTTTGTCTCT CTCGACTGATCTAACAATCC CTAATCTGTGTTCTAAATTC 60
 CTGGACGAGATTTGACAAAG TCCGTATAGCTTAACCTGGT TTAATTTCAAGTGACAGAT 119
 ATG CCC CTT ATT CAT CGG AAA AAG CCG ACG CAG AAA CCA TCG ACG CCG 167
 Met Pro Leu Ile His Arg Lys Lys Pro Thr Glu Lys Pro Ser Thr Pro
 CCA TCT GAA GAG GTG GTG CAC GAT GAG GAT TCG CAA AAG AAA CCA CAC 215
 Pro Ser Glu Glu Val Val His Asp Glu Asp Ser Gln Lys Lys Pro His
 GAA TCT TCC AAA TCC CAC CAT AAG NAA TCG AAC GGA GGA GGG AAG TGG 263
 Glu Ser Ser Lys Ser His His Lys ??? Ser Asn Gly Gly Gly Lys Trp
 TCG TGC ATC GAT TCT TGT TGT TGG TTC ATT GGG TGT GTG TGT GTA ACC 311
 Ser Cys Ile Asp Ser Cys Cys Trp Phe Ile Gly Cys Val Cys Val Thr
 TGG TGG TTT CTT CTC TTC CTT TAC AAC GCA ATG CCT GCG AGC TTC CCT 359
 Trp Trp Phe Leu Leu Phe Leu Tyr Asn Ala Met Pro Ala Ser Phe Pro
 CAG TAT GTA ACG GAG CCG AAT CAC GNG TCC TTT GCC TTA CCC G 402
 Gln Tyr Val Thr Glu Pro Asn His ??? Ser Phe Ala Leu Pro

00000-00000

<210> 96
 <211> 1562
 <212> genomic DNA
 <213> Arabidopsis thaliana
 <400> 9

ATGAAAAAAA TATCTTCACA TTATTCGGTA GTCATAGCGA TACTCGTTGT, 50
 GGTGACGATG ACCTCGATGT GTCAAGCTGT GGGTAGCAAC GTGTACCCTT 100
 TGATTCTGGT TCCAGGAAAC GGAGGTAACC AGCTAGAGGT ACGGCTGGAC 150
 AGAGAATACA AGCCAAGTAG TGTCTGGTGT AGCAGCTGGT TATATCCGAT 200
 TCATAAGAAG AGTGGTGGAT GGTTTAGGCT ATGGTTTCGAT GCAGCAGTGT 250
 TATTGTCTCC CTCACCAGG TGCTTCAGCG ATCGAATGAT GTTGTAATAT 300
 GACCTGTATT TGGATGATTA CCAAAATGCT CCTGGTGTCC AAACCCGGGT 350
 TCCTCATTTC GGTTCGACCA AATCACTTCT ATACCTCGAC CCTCGTCTCC 400
 GGTTAGTACT TTCCAAGATA TATCATTTTG GGACATTTGC ATAATGAACA 450
 AAATAGACAT AAATTTGGGG GATTATTGTT ATATCAATAT CCATTTATAT 500
 GCTAGTCGGT AATGTGAGTG TTATGTTAGT ATAGTTAATG TGAGTGTAT 550
 GTGATTTTCC ATTTTAAATG AAGCTAGAAA GTTGTCGTTT AATAATGTTG 600
 CTATGTCATG AGAATTATAA GGACACTATG TAAATGTAGC TTAATAATAA 650
 GGTTTGATTT GCAGAGATGC CACATCTTAC ATGGAACATT TGGTGAAAGC 700
 TCTAGAGAAA AAATGCGGGT ATGTTAACGA CCAAAACCATC CTAGGAGCTC 750
 CATATGATTT CAGGTACGGC CTGGCTGCTT CGGGCCACCC GTCCCGTGTA 800
 GCCTCACAGT TCCTACAAGA CCTCAAACAA TTGGTGGAAA AACTAGCAG 850
 CGAGAACGAA GGAAAGCCAG TGATACTCCT CTCCCATAGC CTAGGAGGAC 900
 TTTTCGTCCT CCATTTCTTC AACCGTACCA CCCCTTCATG GCGCCGCAAG 950
 TACATCAAAC ACTTTGTTGC ACTCGCTGCG CCATGGGGTG GGACGATCTC 1000
 TCAGATGAAG ACATTTGCTT CTGGCAACAC ACTCGGTGTC CCTTTAGTTA 1050
 ACCCTTTGCT GGTGAGACGG CATCAGAGGA CCTCCGAGAG TAACCAATGG 1100
 CTACTTCCAT CTACCAAAGT GTTTCACGAC AGAACTAAAC CGCTTGTCGT 1150
 AACTCCCCAG GTTAACTACA CAGCTTACGA GATGGATCGG TTTTTTGACG 1200
 ACATTGGATT CTCACAAGGA GTTGTCCTT ACAAGACAAG AGTGTGTCCT 1250

00537740.033000

TTAALCAGAGG	AGCTGATGAC	TCCGGGAGTG	CCAGTCACTT	GCATATATGG	1300
GAGAGGAGTT	GATACACCGG	AGGTTTTTGAT	GTATGGAAAA	GGAGGATTCG	1350
ATAAGCAACC	AGAGATTAAAG	TATGGAGATG	GAGATGGGAC	GGTTAATTTG	1400
GCGAGCTTAG	CAGCTTTGAA	AGTCGATAGC	TTGAACACCG	TAGAGATTGA	1450
TGGAGTTTCG	CATACATCTA	TACTTAAAGA	CGAGATCGCA	CTTAAAGAGA	1500
TTATGAAGCA	GATTTCAATT	ATTAATTATG	AATTAGCCAA	TGTTAATGCC	1550
GTCAATGAAT	GA				1562

<210> 10
 <211> 3896
 <212> genomic DNA
 <213> Arabidopsis thaliana
 <400> 10

ATGGGAGCGA	ATTGAAATC	AGTAACGGCT	TCCTTCACCG	TCATCGCCGT	50
TTTTTCTTG	ATTTGCGGTG	GCCGAACCTGC	GGTGGAGGAT	GAGACCGAGT	100
TTCACGGCGA	CTACTCGAAG	CTATCGGGTA	TAATCATTCG	CCCAATTTCG	150
TCGACGCAGC	TACGAGCGTG	GTCGATCCTT	GACTGTCCAT	ACACTCCGTT	200
CGACTTCAAT	CCGCTCGACC	TCGTATGGCT	AGACACCACT	AAGGTCCGTTG	250
ATCTTCATTT	CCTTCGCTCC	TTATTCTGTC	GGTCGAGTCA	CTTGTTGATG	300
AATTCCAAGC	GAAATATAGC	AATGAAGCAT	GTCTCGTCTC	TCATTATTGAT	350
TCGTTTATTA	GTCAACAGTG	ACGCTTCTGA	ATCTGAGTTT	AGAGTCATAT	400
AAAACAGCTG	ACTCGGCGAG	TCTTTCCCAT	CGCTTTTGGT	TCGCTAAATG	450
TAGCGCAATG	AATGTGTAAT	TAGTCTGCGC	TTTTTATTCA	ACTAGATCTG	500
CAAGTTTTTC	AGAGTGCTCA	ATAGTAGTTA	GAAAATGTTA	GGTCATTTTA	550
CTTGTTGCAAT	GTGATTCTTT	TGGTTGTTGC	TTACTGATCG	ACGTGATGGA	600
TGGTTTACAG	CTTCTTTCTG	CTGTCAACTG	CTGGTTTAA	TGTATGGTGC	650
TAGATCCTTA	TAATCAAACA	GACCATCCCG	AGTGTAAGTC	ACGGCCTGAC	700
AGTGGTCTTT	CAGCCATCAC	AGAATTGGAT	CCAGGTTACA	TAACAGGTAG	750
TTTCGGATTT	TTCTTTCTTT	TGAGTTTTCT	TCAATTTGAT	ATCATCTTGT	800
TGTGATATAA	TATGGCTAAG	TTCATTAATT	TGGTCAATTT	TCAGGTCCCTC	850
TTTCTACTGT	CTGGAAGAG	TGGCTTAAGT	GGTGTGTTGA	GTTTGGTATA	900
GAAGCAATG	CAATTGTGCG	TGTTCCATAC	GATTGGAGAT	TGTCACCAAC	950
CAATTGGAA	GAAGGTGACC	TTTACTTTCA	CAAGCTCAAG	TTAGTCCTTA	1000
TCAGGCTAAT	GTCTTTTATC	TTCTCTTTT	ATGTAAGATA	AGCTAAGAGC	1050
TCTGGTCGTC	TTCTTTTTTG	CAGGTTGACC	TTTGAAGCTG	CTTTAAAGCT	1100
CCGTGGCGGC	CCTTCTATAG	TATTTGCCCA	TTCAATGGGT	AATAATGTCT	1150
TCAGATACTT	TCTGGAATGG	CTGAGGCTAG	AAATTGCACC	AAAACATTAT	1200
TTGAAGTGCC	TTGATCAGCA	TATCCATGCT	TATTTGCTGT	TGGTACCCGG	1250
CCTACTATCC	TTAAGTTACC	ATTTTATTTT	TTCTCTAATT	CGGGGAGTTA	1300
TGTTGTGACT	TACTGGATTG	AGCTCGATAC	CTGATTTGTT	GTTGATTTAG	1350
GAGCTCCTCT	TCTTGGTTCT	GTTGAGGCAA	TCAAACTCTAC	TCTCTCTGGT	1400
GTAACGTTTG	GCCTTCCTGT	TTCTGAGGTG	ACCTCTGACT	TCTCTTTAGT	1450
TTTAAGTAGT	TGATATCAAC	CAGGTCTTAT	AACTCACTGG	ATTTTCCTTT	1500
TGAAGTAGTT	ACTTTTGTTA	ATTGAACCTG	TGTACGCGAT	ATGGTATCTG	1550
TAGATCTTGA	AGTGTAGTTT	ATCAAAGAAC	ATATTGTGGG	TAGTATACCT	1600
GTCAGCGGCC	TTAGCTAATA	CAACCAAACC	ACATGTACAC	TGATTTAGTT	1650
TTGAGATTAT	TATGGTAGAC	TTTAAGTTGA	GAAGAACTT	TGACTGAAT	1700
CTTTTATATT	TAATAGGCTA	TGATTTGTTT	ATTGAATCA	TGTGACATAT	1750
TGACATGCGC	TTCTCATGTT	TTTTGTTGGC	AAGGCTTCAG	GGAACTGCTC	1800
GGTTGTTGTC	CAATTCTTTT	GCGTCGTCAT	TGTGGCTTAT	GCCATTTTCA	1850
AAGAATTGCA	AGGGTGATAA	CACATTCTGG	ACGCATTTTT	CTGGGGGTGC	1900
TGCAAGAAA	GATAAGCGCG	TATACCACTG	TGATGAAGAG	GAATATCAAT	1950
CAAAATATTC	TGGCTGGCCG	ACAAATATTA	TAAACATTGA	AATTCCTTCC	200
ACTAGCGGTT	AGACTCTGTA	TATGCAACTG	TAACTAATAC	AAAAGTTTCA	2050
CCAAGAATGT	TCACTCTCAT	ATTTCTCTCC	TTTGATGTGT	ATCCATCAGT	2100
TACAGAAACA	GCTCTAGTCA	ACATGACCAG	CATGGAATGT	GGCCTTCCCA	2150
CCCTTTTGTG	TTTCAAGGCC	CGTGAAGTAC	CAGATGGGAC	TCTTTTCAAA	2200
GCAATAGAA	ACTATGACCC	AGATAGCAAG	AGGATGTTAC	ACCAGTTAAA	2250
GAAGTACGTA	CCTTTCTTTG	TGATAAGAAA	TATTGCTCAT	CGATCATCAC	2300
TGCTGGCTT	CTTGACGTC	AAATTGTTTT	GTTTAAATCT	CTATATCAAT	2350
TGTTTATATG	CTTTGTCTTT	CTTACTATAA	GAAACAAGTA	TAATCAGAAA	2400
CCTTATTATT	GATTATCAGT	TCTCTCTTTA	TATTATGGAA	TGTCTTTTTC	2450
GTITACAGTT	ATGAATGCAA	AAGGGGGTAT	TTTAGTTGAT	TGATTCTCTC	2500
ATTCTCTAGT	TTGTTTTGAC	TAATAGCGTC	AATTTTGTGT	TTCTAGCAAA	2550
TCTTTGTCAA	TTATATATAA	CATGCTAAT	ATACTTTTCA	GGTTGTATCA	2600
TGATGACCC	GTTTTTAATC	CTCTGACTCC	TTGGGAGAGA	CCACCTATAA	2650
AAATGTATT	TTGCATATAT	GGTGCTCATC	TAAAGACAGA	GGTATGATGC	2700
ATTCTCAATA	TCACATTATG	CGTTGACTTT	GTTATTATAT	TCCCCATTGT	2750

00537710.030000


2800
2850
2900
2950
3000
3050
3100
3150
3200
3250
3300
3350
3400
3450
3500
3550
3600
3650
3700
3750
3800
3850
3896

<210> 11b
 <211> 709
 <212> cDNA
 <213> tomato
 <400> 11

CTGGGGCCAA AAGTGAACAT AACAAGGACA CCACAGTCAG AGCATGATGT	50
TCAGATGTAC AAGTGCATCT AAATATAGAG CATCAACATG GTGAAGATAT	100
CATTCCCAAT ATGACAAAGT TACCTACAAT GAAGTACATA ACCTATTATG	150
AGGATTCTCA AAGTTTTCCA GGGACAAGAA CAGCAGTTTG GGAGCTTGAT	200
AAAGCAAATC ACAGGAACAT TGTCAGATCT CCACCTTTGA TCGGGGAGCT	250
GTGGCTTGAG ATGTGGCATG ATATTCATCC TGATAAAAAG TCCAAGTTTG	300
TTACAAAAGG TGGTGTCTGA TCCTCACTAT TTTCTTCTAT AAATGTTTGA	350
GTITGTATTG ACATTGTAAG TATTGCAACA AAAAGCAAAG CGTGGGCCTC	400
TGAGGGATCA GGACTGCTAT TGGGATTACG GGAAAGCTCG ATGTGCATGG	450
GCTGAACATT GTGAATACAG GTTAGAATAT TCAAATTATA TTTTGCAAAA	500
TATTCTCTTT TTGTGTATTT AGGCCACCTT TCCCCGGTCA CAACGATGCA	550
GATATGTATT CGGGGATGTT CACCTGGGAC AGAGTTGCAG ATTGAAGAGT	600
TCTACATCTC ACATCCTGTC ACACTATGTG TGATATTTAA GAACTTTGT	650
TTGGCGGAAC AACAAAGTTTG CACAAACATT TGAAGAAGAA AGCGAAATGA	700
TTGAGAGAG	709

0053740.030000

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies were found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☒ Scanned copy is best available.

The first six pages of the sequence listing are the drawings.